

Stressed pairing in conventional color superconductors is unavoidable

Krishna Rajagopal^{1,2,*} and Andreas Schmitt^{1,†}

¹*Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139*

²*Nuclear Science Division, MS 70R319, Lawrence Berkeley National Laboratory, Berkeley, CA 94720*

(Dated: February 2, 2008)

At sufficiently high densities, cold dense three-flavor quark matter is in the color-flavor locked (CFL) phase, in which all nine quarks pair in a particularly symmetric fashion. Once the heaviness of the strange quark (mass m_s) and the requirements of electric and color neutrality are taken into account, the CFL pattern of color superconductivity requires the pairing of quarks that would, in the absence of pairing, have Fermi momenta that differ by of order m_s^2/μ , with μ the quark number chemical potential. This means that at sufficiently small μ , the “stress” on the pairing is large enough that the system can lower its energy by breaking pairs, resulting in some unconventional color superconductor which includes gapless excitations, spatial inhomogeneity, counter-propagating currents, or all three. In this paper we ask whether there is some less symmetric but still conventional pattern of pairing that can evade the stress. In other words, is there a pattern of pairing in which, once electric and color neutrality are imposed by suitable chemical potentials, pairing only occurs among those quarks whose Fermi momenta would be equal in the absence of pairing? We use graph-theoretical methods to classify 511 patterns of conventional color superconducting pairing, and show that none of them meet this requirement. All feel a stress, and all can be expected to become unstable to gapless modes at a density comparable to that at which the CFL phase becomes unstable.

PACS numbers: 12.38.Mh, 24.85.+p

I. INTRODUCTION

A. Context

At any densities that are high enough that nucleons are crushed into quark matter, the quark matter that results must, at sufficiently low temperatures, be in one of a family of color superconducting phases [1]. The essence of color superconductivity is quark pairing, driven by the BCS mechanism which operates whenever there is an attractive interaction between fermions at a Fermi surface [2]. The QCD quark-quark interaction is strong and is attractive between quarks that are antisymmetric in color, so we expect cold dense quark matter to generically exhibit color superconductivity. Color superconducting quark matter may well occur in the cores of compact stars. Except during the first few seconds after their birth, compact stars are very cold compared to the 10’s of MeV critical temperatures expected for color superconductivity, and we shall therefore work at zero temperature throughout this paper.

It is by now well-established that at asymptotic densities, where the up, down and strange quarks can be treated on an equal footing and the disruptive effects of the strange quark mass can be neglected, quark matter is in the color-flavor locked (CFL) phase, in which quarks of all three colors and all three flavors form Cooper pairs [1, 3]. The CFL phase is a color superconductor

but is an electromagnetic insulator, with zero electron density. To describe quark matter as may exist in the cores of compact stars, we need consider quark number chemical potentials μ of order 500 MeV at most (corresponding to baryon number chemical potentials $\mu_B = 3\mu$ of order 1500 MeV at most). The relevant range of μ is therefore low enough that the strange quark mass m_s cannot be neglected: it is expected to be density dependent, lying between the current mass ~ 100 MeV and the vacuum constituent quark mass ~ 500 MeV. In bulk matter, as is relevant for compact stars where we are interested in kilometer-scale volumes, we must furthermore require electromagnetic and color neutrality [4, 5, 6, 7, 8] (possibly via mixing of oppositely charged phases) and allow for equilibration under the weak interactions. All these factors work to pull apart the Fermi momenta of the cross-species pairing that characterizes color-flavor locking. At the highest densities, we expect CFL pairing, but as the density decreases the combination of nonzero m_s and the constraints of neutrality put greater and greater stress on cross-species pairing, and reduce the excitation energies of those fermionic quasiparticles whose excitation would serve to ease the stress by breaking pairs [9].

If we imagine beginning in the CFL phase at asymptotic density, reducing the density, and suppose that CFL pairing is disrupted by the heaviness of the strange quark before color superconducting quark matter is superseded by the hadronic phase, the CFL phase must be replaced by some phase of quark matter in which there is less, and less symmetric, pairing. The nature of this next phase down in density is currently not established.

Within a spatially homogeneous ansatz, the next phase down in density is the gapless CFL (gCFL) phase [9, 10, 11, 12, 13, 14, 15, 16]. In this phase, quarks of all three

*Electronic address: krishna@lns.mit.edu

†Electronic address: aschmitt@lns.mit.edu

colors and all three flavors still form Cooper pairs, but there are regions of momentum space in which certain quarks do not succeed in pairing, and these regions are bounded by momenta at which certain fermionic quasiparticles are gapless. These gapless quasiparticles (and a nonzero density of electrons) make the gCFL phase an electric conductor. This variation on BCS pairing — in which the same species of fermions that pair feature gapless quasiparticles — was first proposed for two flavor quark matter [17] and in an atomic physics context [18]. In all these contexts, however, the gapless paired state turns out in general to be unstable: it can lower its energy by the formation of counter-propagating currents [19, 20]. In the atomic physics context, the resolution of the instability is phase separation, into macroscopic regions of two phases in one of which standard BCS pairing occurs and in the other of which no pairing occurs [21]. This picture appears to be supported by very recent experiments [22, 23]. In the QCD context, the analogue of this phase separation would require coexisting phases one of which is positively charged and the other of which is negatively charged. This may in fact be the resolution to the instability of gapless color superconductivity in two-flavor quark matter [24]. In three-flavor quark matter, where the instability of the gCFL phase has been established in Refs. [20], phase coexistence would require coexisting components with opposite color charges, in addition to opposite electric charges, making it very unlikely that a phase separated solution can have lower energy than the gCFL phase [10, 11]. The gluon condensate phase of Ref. [25] is another possible resolution to the instability in two-flavor quark matter, but it requires an unbroken color $SU(2)$. In the three-flavor quark matter context, it seems likely that a ground state with counter-propagating currents is required. This could take the form of a crystalline color superconductor [26, 27, 28] — the QCD analogue of a form of non-BCS pairing first considered by Larkin, Ovchinnikov, Fulde and Ferrell [29]. Or, given that the CFL phase itself is likely augmented by kaon condensation [30], it could take the form of a phase in which a CFL kaon condensate carries a current in one direction balanced by a counter-propagating current in the opposite direction carried by gapless quark quasiparticles [31]. This meson supercurrent phase has been shown to have a lower free energy than the gCFL phase. The investigation of crystalline color superconductivity in three-flavor QCD has only just begun [28], and it remains to be seen whether such a phase can have a lower free energy than the meson current phase, making it a possible resolution to the gCFL instability. The simplest “crystal” structures do not suffice, but experience in the two-flavor context [27] suggests that realistic crystal structures constructed from more plane waves will prove to be qualitatively more robust.

B. Purpose

The analysis of unconventional pairing in color superconductivity is challenging, given the myriad non-BCS possibilities. In this context, our purpose in this paper is to make an exhaustive search through all the possibilities for conventional BCS pairing in three-flavor quark matter that involve pairing patterns that are less symmetric than CFL. The first and best studied of these is the 2SC phase [32, 33, 34], in which only quarks of two colors and two flavors pair. There are, however, many other possibilities. The question we ask in this paper is whether any of these patterns of pairing allow conventional BCS pairing to persist down to a value of the μ that is parametrically lower than that at which the CFL phase breaks down. We shall find that no such phases exist.

We shall only consider patterns of pairing in which the Cooper pairs are antisymmetric in color, as that is the channel in which the interaction between quarks is attractive. Furthermore, we focus on $J = 0$ pairing (J being the total angular momentum of a Cooper pair) since $J = 1$ pairing between quarks of the same flavor yields gaps that are several orders of magnitude smaller [33, 35, 36]. Antisymmetry in color, together with $J = 0$, requires antisymmetry in flavor. We shall therefore consider all patterns of BCS pairing that are antisymmetric in both color and flavor.

The logic of our investigation is perhaps best explained with reference to the CFL phase itself, see Fig. 1, although the point of our investigation is to apply this logic to other phases. In the CFL phase, color neutrality requires a nonzero chemical potential μ_8 for the color charge proportional to T_8 given by $\mu_8 = -m_s^2/(2\mu)$ [6, 7], whereas electric neutrality requires $\mu_e = 0$ [37]. In the presence of these chemical potentials, but in the absence of pairing, the Fermi momenta of the nine quark species would be given by the effective chemical potentials plotted in Fig. 1. The merit of this figure is that one can immediately see that the pattern of pairing requires pairing between species with differing effective chemical potentials, separated by $\delta\mu_{\text{eff}} \sim m_s^2/\mu$. In this circumstance, the BCS state describing, for example, gs - bd pairing is constructed by first filling the gs and bd states up to a common Fermi surface given by the average between their two effective chemical potentials — which exacts a free energy price of order $\delta\mu_{\text{eff}}^2 \sim m_s^4$ — and then pairing — which gains a free energy benefit of order $\Delta^2 \mu^2$. In the CFL phase at zero temperature the gap parameter Δ is estimated to be of order 10 to 100 MeV [1, 3]. When the density drops sufficiently that the quark chemical potential μ falls below some critical value of order m_s^2/Δ , then, the cost of pairing must exceed the benefit gained from pairing, gs - bd pairs must break, and the gCFL phase at that point takes over from the CFL phase. At the CFL-gCFL transition, however, these two phases have the same free energy. This means that given that the gCFL phase is itself unstable, there must be some other




FIG. 1: Schematic diagram of the effective chemical potentials μ_{eff} for the nine quarks as a function of the strange mass, in the CFL phase. We abbreviate red, green and blue by r , g and b and up, down and strange by u , d and s . (The effective chemical potentials are best thought of as the Fermi momenta that the quarks would have in the absence of pairing but in the presence of the chemical potentials μ_e , μ_3 and μ_8 that serve to maintain electric and color neutrality in the paired state.) The splitting between effective chemical potentials is proportional to m_s^2/μ . The quarks encircled with the symbol of the same shape (and color on line) pair with each other in the CFL phase. We see two instances of stressed pairing — i.e. pairing between quarks with differing effective chemical potentials. Such pairing gains a free energy benefit of order $\Delta^2\mu^2$ but also exacts a free energy cost of order $(m_s^2/\mu)^2\mu^2 = m_s^4$. Within a spatially homogeneous ansatz, the CFL phase is superseded by the gCFL phase — in which some bu - rs and gs - bd pairs break — when $m_s^2/\mu > 2\Delta$. However, the gCFL phase is itself unstable, meaning that the CFL phase must in fact be superseded earlier by some other phase. In this paper we ask whether any conventional BCS-paired state involves pairing only between quarks with the same effective chemical potential. Such a phase could take over from the CFL phase and persist down to a density for which $m_s^2/\mu \gg \Delta$. We find no such phase.

phase which takes over from CFL before the CFL-gCFL transition is reached. In this investigation, we suppose that this next-to-highest density phase of quark matter involves conventional BCS pairing only, and involves pairing that is antisymmetric in color and flavor only. We then ask whether there is any pattern of such pairing in which only quarks with the same effective chemical potential pair.

If the answer to this question were yes, we would then have a candidate pattern of pairing (or several candidate patterns) which could persist down to much lower densities than the CFL phase, where $m_s^2/\mu \gg \Delta$, without be-

coming unstable towards pair-breaking, and hence without the need to face the question of what unconventional non-BCS state resolves the instability of a gapless color superconducting state towards the nucleation of counter-propagating currents. We would then have to derive and solve the gap equation for this stress-free pattern of pairing, studying it more fully than in the analysis we present here.

In fact, we find no patterns of stress-free pairing. Among all the patterns of color and flavor antisymmetric BCS pairing that are possible in three-flavor quark matter, stressed pairing is unavoidable. We conclude that even if the CFL phase were supplanted by some less symmetric pattern of otherwise still conventional BCS pairing, any such phase would itself become unstable to pair breaking for some m_s^2/μ of order Δ . This means that unless Δ/m_s^2 is large enough that the CFL phase persists down to low enough densities that it is superseded by hadronic matter, analysis of non-BCS options, including spatial inhomogeneity and/or counter-propagating currents and/or gapless color superconductivity, is unavoidable.

The remainder of the paper is organized as follows. In Sec. II, we present a method to implement neutrality conditions in superconducting matter for a generic pattern of pairing. This method is a generalization of that applied to the CFL and 2SC phases in Ref. [6]. We shall assume that m_s/μ is small, but as Fig. 1 makes clear this is sufficient to answer the question of whether stress-free pairing is possible. In Sec. III, we present two simple toy models with two and three fermion species, respectively, in order to illustrate the interplay between the pairing pattern and the neutrality conditions. In Sec. IV we describe the application of our method to three-flavor quark matter, including subtleties associated with imposing color neutrality. Some of these subtleties are explained further in Appendix B. Sec. V contains the central argument and results of the paper.

II. NEUTRALITY CONDITIONS AND STRESS-FREE STATES

In this Section, we present a general form of the neutrality conditions in a superconducting state, to leading order in small parameters in a sense that we shall explain. In subsequent sections, we shall use this formalism first to analyze two simple toy models and then to analyze the multitude of pairing patterns that are possible in quark matter with three colors and three flavors.

Consider a system of N fermions with a common chemical potential μ and K different “charge chemical potentials” μ_{Q_k} , meaning that the N fermions have chemical potentials

$$\mu_i = \mu + \sum_{k=1}^K q_{ik} \mu_{Q_k}, \quad i = 1, \dots, N. \quad (1)$$

The coefficient q_{ik} is the Q_k -charge of the i -th fermion. Neutrality with respect to K different Q_k -charges then requires that the free energy Ω satisfy

$$\frac{\partial \Omega}{\partial \mu_{Q_k}} = 0, \quad k = 1, \dots, K. \quad (2)$$

In Sections IV and V, where we shall apply this formalism to quark matter, N will be 9 (quarks of three colors and three flavors) and K will be 3, because electric and color neutrality are maintained via the adjustment of three chemical potentials μ_e , μ_3 and μ_8 coupling to electric, color- T_3 , and color- T_8 charges. In QCD, μ_e , μ_3 and μ_8 are the zeroth components of electromagnetic and color gauge fields, and the gauge field dynamics ensure that they take on values such that the matter is neutral, satisfying (2) [6, 38, 39].

We shall only consider charges Q_k which are traceless,

$$\sum_i q_{ik} = 0, \quad (3)$$

as this suffices for the analysis of three-flavor quark matter. If all the charges with respect to which neutrality must be maintained are traceless, as we assume, then if all the fermions have the same mass in the absence of pairing the matter is neutral with all μ_{Q_k} set to zero. We shall assume, however, that some of the fermions have mass m while others are massless. (In three-flavor quark matter m will be the strange quark mass, and we shall treat the light quarks as massless.) We shall always assume that m and all the μ_{Q_k} are much smaller than the common μ . We shall then see that neutrality requires that the μ_{Q_k} are parametrically of order m^2/μ .

Our analysis can be applied to three-flavor quark matter in which not all quarks pair, but it cannot be applied directly to two-flavor quark matter since in that context electric charge is not traceless. However, the question that we address in three-flavor quark matter in this paper can be answered with much less effort in the two-flavor case, in which μ_e is parametrically of order μ in the absence of pairing. This makes it easy to show that there are no stress-free patterns of pairing in two-flavor quark matter that are antisymmetric in color and flavor.

We shall consider many different patterns of pairing, but in each case we shall simplify our notation by pretending that any quarks which pair do so with a common gap parameter Δ , assumed to be $\ll \mu$. In any realistic context, the pattern of pairing obtained by solving some gap equation may well have different gap parameters for different pairs of quarks. In such a case, the Δ appearing in our analysis should be taken to be the smallest nonzero gap parameter. We shall also see, momentarily, that the magnitude of the gap parameter Δ does not actually play a role in our analysis; all that matters to us is the pattern of which quarks pair.

The free energy of a conventional BCS state can be described as if one first assigns a common Fermi momentum

ν to all the fermions that pair with each other, and then reduces the free energy relative to that of this fictional unpaired state by a condensation energy of order $\Delta^2 \nu^2$. If the “free energy benefit” of pairing (of order $\Delta^2 \nu^2$) is larger than the “free energy cost” of first equalizing the Fermi momenta, then the BCS state under consideration is stable against pair-breaking. In this paper, we shall seek “stress-free states”, for which the pattern of pairing is chosen such that the free energy cost of equalizing the Fermi momenta is zero.

As one can see from Fig. 1, the CFL state has four sets of quarks that are “connected” by its specific pattern of pairing, meaning that in its analysis we will have to introduce four different common Fermi momenta ν_n . Anticipating the terminology from graph theory that we shall use in Sec. IV, we term these sets of fermions *components*. The components of the CFL pattern are $\{bu, rs\}$, $\{bd, gs\}$, $\{rd, gu\}$, and $\{ru, gd, bs\}$. Unpaired quark matter has nine components, each consisting of a single quark. For the general case, we denote the components by $\mathcal{C}_n$, with $n = 1 \dots c$ where c denotes the number of components. Thus, $c = 4$ for the CFL pattern. We shall assume that the n -th component $\mathcal{C}_n$ contains r_n massless fermions and s_n fermions with mass m . We now give an expression for the free energy Ω , which generalizes that in Ref. [6], assuming that fermions in components $\mathcal{C}_n$ pair with gap parameter Δ :

$$\Omega = \frac{1}{\pi^2} \sum_{n=1}^c \int_0^{\nu_n} dp p^2 \left(r_n p + s_n \sqrt{p^2 + m^2} - \sum_{i=1}^{r_n+s_n} \mu_i \right) - \mathcal{N} \frac{\Delta^2 \mu^2}{4\pi^2}. \quad (4)$$

We have denoted the common Fermi momenta by ν_n ; these are best thought of as variational parameters and will be determined below by minimizing Ω with respect to them. The condensation energy is written somewhat schematically, as it is of no consequence in our analysis as we now explain. As long as $\Delta \ll \mu$, as we assume, we can neglect terms of order $\Delta^2 \mu \mu_{Q_k}$ and $\Delta^2 \mu_{Q_k}^2$ in the condensation energy, as they are much smaller than the $\mu^2 \mu_{Q_k}^2$ terms that will turn out to determine the free energy cost of equalizing Fermi momenta at the values ν_i . Once the $\Delta^2 \mu_{Q_k}^2$ terms are dropped, the condensation energy is as written, with $\mathcal{N}$ a number that depends on the pattern of pairing. ($\mathcal{N}$ counts the number of pairs of fermions that pair; $\mathcal{N} = 12$ in the CFL pattern, for example.) Upon making this approximation the condensation energy is independent of the charge chemical potentials and hence irrelevant in the neutrality conditions. We therefore neglect it henceforth. Another way of saying this is that our purpose is to seek patterns of pairing that minimize the free energy cost of equalizing the Fermi momenta of those quarks that pair. If we can eliminate this cost, we will have stress-free pairing. As we will not be evaluating the free-energy benefit of pairing, the detailed form of the condensation energy is of no consequence.

We now minimize Ω with respect to the c different com-

mon Fermi momenta ν_n , one each for the paired fermions in each of the c different components $\mathcal{C}_n$. In so doing, we determine the values of the ν_n in terms of the chemical potentials and m . Because it is clear from Eq. (4) that the equations $\partial\Omega/\partial\nu_n = 0$ are not coupled, we can minimize Ω with respect to each ν_n separately, yielding the condition

$$\nu_n^2 \left(r_n \nu_n + s_n \sqrt{\nu_n^2 + m^2} - \sum_{i=1}^{r_n+s_n} \mu_i \right) = 0 \quad (5)$$

that must be satisfied by each of the ν_n . The nontrivial solution to this condition is

$$\nu_n = \frac{r_n}{r_n^2 - s_n^2} \sum_{i=1}^{r_n+s_n} \mu_i - \frac{s_n}{r_n^2 - s_n^2} \sqrt{\left(\sum_{i=1}^{r_n+s_n} \mu_i \right)^2 + (r_n^2 - s_n^2) m^2} . \quad (6)$$

We now use Eq. (1), and expand this expression for ν_n to linear order in the charge chemical potentials μ_{Q_k} and to quadratic order in m . If, upon working at lowest nontrivial order in these small quantities, we were to find a stress-free pattern of pairing, we would then have to return to this point in the analysis and investigate higher order corrections. As we shall show that there are no stress-free patterns to this order, we shall not need to consider higher order terms. Upon making this approximation, the result for ν_n is most conveniently written in terms of “effective chemical potentials” μ_i^{eff} for each of the fermions which, following Ref. [6], we define as

$$\begin{aligned} \mu_i^{\text{eff}} &\equiv \mu_i && \text{for the massless fermions} \\ \mu_i^{\text{eff}} &\equiv \mu_i - \frac{m^2}{2\mu} && \text{for the fermions with mass } m . \end{aligned} \quad (7)$$

We find

$$\nu_n = \frac{1}{r_n + s_n} \sum_{i=1}^{r_n+s_n} \mu_i^{\text{eff}} . \quad (8)$$

We see that, to the order at which we are working, the common Fermi momentum ν_n of the fermions in the component $\mathcal{C}_n$ is given by the arithmetic mean of the effective chemical potentials of all the fermions in $\mathcal{C}_n$. This result generalizes the one of Ref. [6]. It is also straightforward, although somewhat less instructive, to write the expression for ν_n without introducing effective chemical potentials:

$$\nu_n = \mu + \frac{1}{r_n + s_n} \sum_{i=1}^{r_n+s_n} \sum_{k=1}^K q_{ik} \mu_{Q_k} - \frac{s_n}{r_n + s_n} \frac{m^2}{2\mu} . \quad (9)$$

We now evaluate the neutrality conditions (2). The straightforward but lengthy way to do this is to perform the p -integral in Eq. (4), insert the result (9) for

ν_n into the resulting expression, and differentiate with respect to μ_{Q_k} . Both the calculation and the form of the result are, however, simpler if we first make the following observation. Consider Ω as a function of $\nu_1, \dots, \nu_c$ and $\mu_{Q_1}, \dots, \mu_{Q_K}$, where each ν_n itself is a function of $\mu_{Q_1}, \dots, \mu_{Q_K}$. Then, the neutrality conditions can be written as

$$0 = \frac{d\Omega}{d\mu_{Q_k}} = \sum_{n=1}^c \frac{\partial\Omega}{\partial\nu_n} \frac{\partial\nu_n}{\partial\mu_{Q_k}} + \frac{\partial\Omega}{\partial\mu_{Q_k}} . \quad (10)$$

The first term on the right-hand side of this equation vanishes as long as we are evaluating it at values of the ν_n for which $\partial\Omega/\partial\nu_n = 0$. Consequently, we need only evaluate the second term in which only the explicit μ_{Q_k} -dependence of Ω is relevant. We see from (4) that the μ_{Q_k} occur in Ω only in terms proportional to $\int^{\nu_n} dp p^2 \sim \nu_n^3$, and hence conclude that the neutrality conditions take the form

$$0 = \frac{\partial\Omega}{\partial\mu_{Q_k}} = -\frac{1}{\pi^2} \sum_{n=1}^c \frac{\nu_n^3}{3} \frac{\partial}{\partial\mu_{Q_k}} \sum_{i=1}^{r_n+s_n} \mu_i . \quad (11)$$

We can simplify this upon noting from (1) that

$$\frac{\partial}{\partial\mu_{Q_k}} \sum_{i=1}^{r_n+s_n} \mu_i = \sum_{i=1}^{r_n+s_n} q_{ik} \quad (12)$$

and by rewriting the double sum (first over components, then over fermions within components) as a single sum over all the fermions, assigning the i -th fermion a ν_i given by the ν_n of the component to which it belongs. The resulting neutrality conditions take the form

$$0 = \sum_{i=1}^N \nu_i^3 q_{ik} , \quad k = 1, \dots, K , \quad (13)$$

which is as one might have expected. We can further simplify this result by thinking of ν_i as $\mu - (\mu - \nu_i)$, noting that our analysis is valid only to linear order in m^2 and μ_{Q_k} and hence, according to (9), only to linear order in $(\mu - \nu_i)$, and therefore linearizing ν_i^3 as

$$\nu_i^3 = \mu^3 - 3\mu^2(\mu - \nu_i) = 3\mu^2 \left(\nu_i - \frac{2}{3}\mu \right) . \quad (14)$$

Upon using the traceless condition (3), the neutrality conditions then take on the simple form

$$0 = \sum_{i=1}^N \nu_i q_{ik} . \quad k = 1, \dots, K , \quad (15)$$

In Sections IV and V we shall use the expression (9) for the ν_i 's to apply this condition to a multitude of potential pairing patterns in three-flavor quark matter, after first applying it in a toy-model context in Section III.

With the notation that we have by now introduced, the condition for stress-free pairing can be written simply as

$$\nu_i = \mu_i^{\text{eff}}, \quad i = 1, \dots, N. \quad (16)$$

A stress-free state has the property that the chemical potentials that maintain its neutrality are such that when we write the free energy for the BCS state we discover that those fermions which pair (i.e. those that lie within the same component) already have the same μ^{eff} . Hence, the arithmetic mean in (8) yields simply (16). Our goal, then, is to find a neutral, stress-free, pairing pattern, namely one that simultaneously fulfills Eqs. (15) and (16).

III. TWO TOY MODELS

In this Section, we illustrate the interplay between the neutrality conditions and the pairing pattern in two toy models. The first model consists of two fermion species, one of which is massive. In the second model, we add a third, massless, fermion to this system in order to allow for more pairing patterns.

In the first model, we have two fermion species — $N = 2$ — which means that the most general charge assignments we can consider given the requirement of tracelessness is simply the case with $K = 1$, and charge assignments $q_1 = q$ and $q_2 = -q$. That is, we have fermions with chemical potentials

$$\mu_1 = \mu + q\mu_q, \quad (17)$$

$$\mu_2 = \mu - q\mu_q. \quad (18)$$

We assume the first (second) fermion to be massless (massive), making the effective chemical potentials

$$\mu_1^{\text{eff}} = \mu_1, \quad (19)$$

$$\mu_2^{\text{eff}} = \mu_2 - \frac{m^2}{2\mu}. \quad (20)$$

There are only two possible patterns: pairing and no pairing. Without pairing, the common Fermi momenta are simply the effective chemical potentials, $\nu_i = \mu_i^{\text{eff}}$ and the neutrality condition (15) yields

$$\mu_q = -\frac{m^2}{4q\mu} \quad (21)$$

This is easy to understand: with $m = 0$, the system is neutral with $\mu_q = 0$. Switching on the mass reduces the effective Fermi surface of the second species reducing its density. Consequently, a nonzero μ_q is required in order to restore neutrality, which here simply means equality of the two Fermi surfaces. The value of μ_q is such that $\mu_1^{\text{eff}} = \mu_2^{\text{eff}} = \mu - m^2/(4\mu)$.

When there is Cooper pairing between the two species, there is a single common Fermi momentum

$$\nu_1 = \nu_2 = \frac{\mu_1^{\text{eff}} + \mu_2^{\text{eff}}}{2} = \mu - \frac{m^2}{4\mu}. \quad (22)$$

In this case, the neutrality condition (15), and indeed even the result (13), are automatically satisfied, for any value of μ_q . In this sense, this toy model is too simple, but let us continue with it a little farther before generalizing it. Since we can achieve neutrality with any μ_q , let us choose it according to (21), which makes the effective potentials for the two fermions equal before pairing, satisfying (16) and rendering the paired state stress-free. The fact that μ_q can be varied around its stress-free value without upsetting neutrality is a sign that the material is an insulator. Varying μ_q away from that in (16) introduces a stress on the pairing. There is then a free-energy cost to equalizing the Fermi surfaces, and the paired state is only favored if its condensation energy is large enough. The paired state will therefore be favored in a band of values of μ_q centered on the stress-free value (16), with the width of this band being controlled by the magnitude of Δ . In our analysis, we are not evaluating the condensation energy and so cannot evaluate the width of the bandgap of this insulating state.

We can augment the toy model slightly by introducing “electrons”, massless fermions which by hypothesis do not pair and which have charge q and chemical potential $\mu_q \ll \mu$. These add a term of order μ_q^3 to the neutrality condition, which is subleading relative to those we have already dropped in writing (15), and so if we were using (15) we would not be able to evaluate the consequences of adding electrons. However, as we have satisfied (13), in this simple model, we are justified in including electrons in the neutrality condition, and thus concluding that once we have introduced electrons into the theory we must set $\mu_q = 0$, keeping the electron density zero, in order to maintain neutrality. We now see that in this toy model augmented with the addition of “electrons”, the stress-free pairing condition $\mu_q = -m^2/(4q\mu)$ and neutrality condition $\mu_q = 0$ cannot both be satisfied. Hence, pairing in neutral matter will of necessity be stressed, and the BCS state will only be favored if Δ is large enough or, equivalently, if for some fixed Δ the stress $|\mu_1^{\text{eff}} - \mu_2^{\text{eff}}|$ is small enough. In fact, in this simple context the quantitative criterion is easily evaluated: breaking a pair costs 2Δ in condensation energy and benefits $|\mu_1^{\text{eff}} - \mu_2^{\text{eff}}| = m^2/(2\mu)$ in reduced stress, and so the paired state is favored as long as $\Delta > m^2/(4\mu)$. This toy model thus proves in many ways to be analogous to the CFL phase, analyzed in Refs. [9, 10, 37].

We now consider a second toy model, in which there are several possible pairing patterns to consider. We add a third fermion species, so now $N = 3$, but we continue to keep $K = 1$. We choose one of the three fermions to be massive, and hence consider a toy model in which the effective chemical potentials are

$$\mu_1^{\text{eff}} = \mu + q_1\mu_q, \quad (23a)$$

$$\mu_2^{\text{eff}} = \mu + q_2\mu_q, \quad (23b)$$

$$\mu_3^{\text{eff}} = \mu + q_3\mu_q - \frac{m^2}{2\mu}, \quad (23c)$$




FIG. 2: Possible patterns of pairing in the toy model with 3 fermion species, shown as graphs. Nonzero gap parameters Δ_i yield edges in the graphs. Below each graph, its components are listed. The four graphs in the second row have identical components and thus are equivalent in our analysis, even though they would not have equivalent condensation energies.

where

$$q_1 + q_2 + q_3 = 0. \quad (24)$$

Which pairing patterns are possible in this model? We consider the gap matrix

$$\mathcal{M} = \begin{pmatrix} 0 & \Delta_1 & \Delta_2 \\ \Delta_1 & 0 & \Delta_3 \\ \Delta_2 & \Delta_3 & 0 \end{pmatrix}, \quad (25)$$

where a nonzero entry in the i -th row and j -th column indicates that fermion i forms Cooper pairs with fermion j . The matrix $\mathcal{M}$ is symmetric because we are assuming that the Cooper pairs have zero angular momentum, and are thus antisymmetric in Dirac indices. We have written the most general possible gap matrix with diagonal entries set to zero. In Section IV, the matrix $\mathcal{M}$ will be a matrix in color and flavor space, and we shall only consider pairing that is antisymmetric in both color and flavor. This means that the diagonal entries of $\mathcal{M}$ will vanish there, so we make this choice here too. We will not be solving a gap equation, and hence will only be interested in which of the Δ 's are nonzero. We shall therefore choose each of Δ_1 , Δ_2 and Δ_3 to be either Δ or zero, yielding $2^3 = 8$ possible patterns of pairing. However, some of these patterns are equivalent in their consequences for our analysis, as we now explain. We can denote each of the 8 possible patterns by a graph, where the vertices of the graph are the fermions and the edges are the nonzero gap parameters. In this simple example of three fermion species, this might seem notationally superfluous but in the case of three-flavor quark matter, with nine fermion species, we shall find that the correspondence between a pairing pattern and a graph provides a useful tool to classify the multitude of possible pairing patterns. For further explanation of the

components	neutrality	stress-free pairing
$\{1\}, \{2\}, \{3\}$	$\mu_q = \frac{q_3}{q_1^2 + q_2^2 + q_3^2} \frac{m^2}{2\mu}$	(no pairing)
$\{1,2\}, \{3\}$	$\mu_q = \frac{1}{3q_3} \frac{m^2}{\mu}$	for any μ_q , if $q_1 = q_2$
$\{1,3\}, \{2\}$	$\mu_q = -\frac{1}{3q_2} \frac{m^2}{2\mu}$	$\mu_q = \frac{1}{q_3 - q_1} \frac{m^2}{2\mu}$
$\{2,3\}, \{1\}$	$\mu_q = -\frac{1}{3q_1} \frac{m^2}{2\mu}$	$\mu_q = \frac{1}{q_3 - q_2} \frac{m^2}{2\mu}$
$\{1,2,3\}$	μ_q arbitrary	if $q_1 = q_2$ and $\mu_q = -\frac{m^2}{6q_1\mu}$

TABLE I: The neutrality condition (15) and the condition for stress-free pairing (16) for a toy model with 3 fermion species.

elementary graph-theoretical terms that we shall use in the following, see for instance Ref. [40]. All 8 graphs are shown in Fig. 2, and we see that the last 4 all have the same components — in this case, actually, the same single component. In our analysis, all that matters is the components since these determine which fermions have a common Fermi momenta. We can therefore consider all the graphs that have the same components as equivalent, even though they would have different condensation energies. This reduces the number of possible patterns that we must consider, in this example from 8 to 5. The number of patterns that we need consider is the number of possible partitions of the set $\{1, 2, 3, \dots, N\}$ into disjoint, non-empty subsets. In combinatorics, this number is called the Bell number B_N , and Fig. 2 shows that $B_3 = 5$. In general, B_N is given by the recursive formula [41]

$$B_N = \sum_{k=0}^{N-1} B_k \binom{N-1}{k}, \quad (26)$$

along with the definition $B_0 \equiv 1$. This yields $B_1 = 1$, $B_2 = 2$, $B_3 = 5$, $B_4 = 15$, $\dots$ $B_9 = 21147$, $\dots$. Returning to our model with $N = 3$, we can now discuss each of the 5 patterns separately, analyzing their neutrality conditions and looking for stress-free patterns of pairing. The results are collected in Table I.

The first row of this table describes the unpaired scenario. In this case, a nonzero μ_q is required, as in the two-species toy model.

The second, third, and fourth rows of the table describe patterns in which two of the fermions form Cooper pairs while the third remains unpaired. In the second row, for example, neutrality is obtained as long as

$$\mu_q = \frac{q_3}{\frac{1}{2}(q_1 + q_2)^2 + q_3^2} \frac{m^2}{2\mu}, \quad (27)$$

which yields the result quoted in the table since $q_1 + q_2 = -q_3$. The analyses of the neutrality conditions in the third and fourth rows are similar, and in each of these cases a particular nonzero μ_q is required. The conditions for stress-free pairing in the second, third and fourth rows of the table follow directly from (16) and (23). For each of the patterns in the second, third and fourth row of the table, the condition that the phase is simultaneously neutral and stress-free turns out to be that this is

possible only if $q_1 = q_2 = -q_3/2$, and that it requires $\mu_q = -m^2/(6q_1\mu)$.

The fifth row of the table describes any of the patterns in which all three fermions pair. (This could be any of four different patterns, as we saw in Fig. 2.) In this case, the system is automatically neutral, with any value of μ_q . However, the pairing is stress-free only if $q_1 = q_2 = -q_3/2$ and $\mu_q = -m^2/(6q_1\mu)$.

Our conclusion for this toy model is that if $q_1 \neq q_2$, neutral stress-free pairing is impossible, whereas if $q_1 = q_2$, we have an excess of possibilities, as *all* the patterns of pairing can be both neutral and stress free. This latter result comes about because if $q_1 = q_2 = -q_3/2$ and $\mu_q = -m^2/(6q_1\mu)$ then the unpaired phase has three equal effective chemical potentials. The case of three-flavor quark matter, to which we turn next, is more analogous to the toy model with $q_1 \neq q_2$.

IV. PAIRING PATTERNS, COLOR SYMMETRY, AND COLOR NEUTRALITY

We now turn to the case of three-flavor quark matter. The neutrality conditions which we introduced in previous sections in a general and abstract way shall now be those required to ensure that the quark matter is electrically and color neutral. We first discuss possible pairing patterns and the role of color and flavor symmetries. We then explain how color neutrality is taken into account. This section provides the basis for Sec. V, where we shall demonstrate that stress-free pairing is impossible in three-flavor quark matter.

A. Gap matrix and color/flavor symmetries

In quark matter with three colors and flavors, nine different quarks are present. Were we to ignore all qual-

itative features of the nature of the interaction between quarks in QCD, we would be forced to consider a general 9×9 gap matrix $\mathcal{M}$. Restricting our attention to the $J = 0$ channel which is antisymmetric in Dirac indices (because $J = 1$ pairing yields much smaller gaps [33, 36]) makes $\mathcal{M}$ a symmetric matrix specified by 45 independent gap parameters which, for our purposes, are either zero or nonzero. This would lead to 2^{45} different gap matrices. However, as before many of these gap matrices would have the same set of components, and the number of patterns that have to be treated separately would be given by the Bell number $B_9 = 21147$, much smaller but still unwieldy.

In QCD the one-gluon exchange interaction between two quarks is attractive in channels which are antisymmetric under the exchange of the color of the quarks. We know that this is the channel in which the dominant attraction between quarks is to be found also by analyzing the nonperturbative instanton-induced interaction between quarks, as well as by the qualitative argument that only by combining two color $[\mathbf{3}]_c$ quarks antisymmetrically into a color $[\mathbf{3}]_c$ diquark do we reduce the color flux. Cooper pairs that are antisymmetric in color and Dirac indices must also be antisymmetric in flavor indices, meaning that we need only consider the antisymmetric flavor channel $[\mathbf{3}]_f$. Therefore, we only consider those 9×9 gap matrices $\mathcal{M}$ in the direct product space $[\mathbf{3}]_c \otimes [\mathbf{3}]_f$, which can be written as

$$\mathcal{M} = \Delta_{ij} J_i \otimes I_j, \quad (28)$$

where the antisymmetric 3×3 matrices $(J_i)_{jk} = -i\epsilon_{ijk}$ and $(I_i)_{jk} = -i\epsilon_{ijk}$ are bases of $[\mathbf{3}]_c$ and $[\mathbf{3}]_f$, respectively. We see that the 9×9 matrix is fully specified by only 9 (not 45) gap parameters, namely the 9 entries in Δ_{ij} . Explicitly,

$$\mathcal{M} = \begin{pmatrix} 0 & 0 & 0 & 0 & -\Delta_{33} & \Delta_{32} & 0 & \Delta_{23} & -\Delta_{22} \\ 0 & 0 & 0 & \Delta_{33} & 0 & -\Delta_{31} & -\Delta_{23} & 0 & \Delta_{21} \\ 0 & 0 & 0 & -\Delta_{32} & \Delta_{31} & 0 & \Delta_{22} & -\Delta_{21} & 0 \\ 0 & \Delta_{33} & -\Delta_{32} & 0 & 0 & 0 & 0 & -\Delta_{13} & \Delta_{12} \\ -\Delta_{33} & 0 & \Delta_{31} & 0 & 0 & 0 & \Delta_{13} & 0 & -\Delta_{11} \\ \Delta_{32} & -\Delta_{31} & 0 & 0 & 0 & 0 & -\Delta_{12} & \Delta_{11} & 0 \\ 0 & -\Delta_{23} & \Delta_{22} & 0 & \Delta_{13} & -\Delta_{12} & 0 & 0 & 0 \\ \Delta_{23} & 0 & -\Delta_{21} & -\Delta_{13} & 0 & \Delta_{11} & 0 & 0 & 0 \\ -\Delta_{22} & \Delta_{21} & 0 & \Delta_{12} & -\Delta_{11} & 0 & 0 & 0 & 0 \end{pmatrix}, \quad (29)$$

with the quarks ordered $ru, rd, rs, gu, gd, gs, bu, bd, bs$. The assumptions we have made, which use only gross qualitative features of QCD, have excluded pairing of the

same quarks, e.g., ru with ru , and also pairing of quarks of the same color, e.g., ru with rd , or with the same flavor, e.g., ru with bu . This is the meaning of the ze-

ros in the matrix $\mathcal{M}$. A pairing pattern now is uniquely defined by the (complex) 3×3 matrix Δ_{ij} . Therefore, within our method, the number of possible patterns is $2^9 = 512$, corresponding to choosing each of the nine entries in Δ_{ij} to be either zero or Δ . One of these, with all entries in Δ_{ij} set to zero, describes unpaired quark matter. The other 511 describe patterns of conventional BCS color superconducting pairing that is antisymmetric in both color and flavor. Note that some of these patterns may describe more than one phase. For example, both the CFL and gCFL phases would be described here as having a Δ_{ij} with nonzero diagonal entries and zero off-diagonal entries, the distinction between whether the three diagonal entries are equal or unequal being unnecessary to the determination of whether the pattern of pairing is stress-free. In our search for stress-free pairing described in Section V it will turn out to be most convenient to consider all 511 patterns even though, as we explain now, not all are physically distinct.

In QCD color symmetry is unbroken, but flavor symmetry is explicitly broken both by quark masses and by electric charge assignments. Let us first consider, however, how our analysis would simplify if both flavor and color symmetries were unbroken. In this circumstance, any two matrices $\mathcal{M}_1, \mathcal{M}_2$ that are connected by a rotation in color-flavor space describe the same phase. A color-flavor rotation $U \otimes V$ where $U \in SU(3)_c$ and $V \in SU(3)_f$ acts on the gap matrix as

$$(U \otimes V) \mathcal{M} = \Delta_{ij} U J_i U^T \otimes V I_j V^T. \quad (30)$$

This rotation can equivalently be applied to the matrix Δ which transforms as

$$(U \otimes V) \Delta = U^T \Delta V, \quad (31)$$

Hence, two matrices Δ_1 and Δ_2 describe the same phase if there exists a rotation $U \otimes V$ such that $\Delta_1 = U^T \Delta_2 V$. However, for any complex matrix Δ_2 there is a choice of unitary matrices U and V such that $U^T \Delta_2 V$ is diagonal [42]. If flavor and color were both unbroken symmetries we would therefore be able to restrict our attention to diagonal matrices Δ , specified by only 3 gap parameters Δ_{11}, Δ_{22} and Δ_{33} . The CFL phase corresponds to the case where these three gap parameters are equal [1, 3]. The gap equations for the more general case where all three can be nonzero and unequal have been derived and solved in Refs. [9, 10, 13], which means that if this were the *most* general case the present analysis would not be necessary.

We must, however, consider the consequences of the explicit breaking of flavor symmetry. In the case $m_u = m_d = 0, m_s \neq 0$, which we shall consider, there appears to be a remaining unbroken $SU(2)_f$ symmetry. However, because the u and d quarks have different electric charges, they will have different effective chemical potentials in the presence of a nonzero chemical potential μ_e as is in general required to impose electric neutrality given the explicit flavor symmetry breaking introduced

by $m_s \neq 0$. We therefore assume that there is no residual $SU(2)_f$ symmetry. Then, two phases Δ_1 and Δ_2 are equivalent only if they are connected by a color rotation, i.e., if there exists a $U \in SU(3)_c$ with $\Delta_1 = U^T \Delta_2$. This means that we must consider both diagonal and non-diagonal matrices Δ . As we explain in the next subsection, this complicates the analysis of the consequences of color neutrality.

As an aside, note that a different analysis is required for the cases of superfluid ${}^3\text{He}$ and for $J = 1$ one-flavor color superconductors, where the relevant symmetry groups are $SO(3)_L \times SO(3)_S$ and $SU(3)_c \times SO(3)_J$, respectively. None of these symmetries are explicitly broken, and yet in both cases, non-diagonal gap matrices must be considered, as for example in the A phase in both systems [43, 44]. The reason is that although a general complex matrix can be diagonalized by left- and right- multiplication by unitary matrices, it cannot be diagonalized if one or other matrix is required to be orthogonal rather than unitary. In Appendix A, we explicitly show that there is no “ A phase” in the case where the unbroken symmetry is $SU(3)_c \times SU(3)_f$. In three-flavor QCD, unlike in the other two cases, non-diagonal matrices need only be considered in the presence of explicit symmetry breaking.

B. Imposing color neutrality

In order to impose color neutrality within our approach, we need to know the corresponding “color charge chemical potentials” μ_{Q_k} , introduced in Eq. (1). Two such chemical potentials are those originating from nonzero expectation values of the zeroth component of the gluon fields A_3^0 and A_8^0 . These chemical potentials (equivalently, gluon fields) couple to the differences between color-charge densities, and so it is no surprise that in QCD their field equations are such as to enforce that they take on values corresponding to the color neutrality conditions (2) [38, 39]. However, we show in Appendix B that, consistent with the results of Ref. [45], in the case of pairing patterns described by non-diagonal matrices Δ , color neutrality may require nonzero expectation values for the zeroth components of *any* of the gluon fields A_a^0 , $a = 1, \dots, 8$. We shall denote these by μ_a , and refer to them as color chemical potentials, even though this terminology is only appropriate for μ_3 and μ_8 . All of them except μ_3 and μ_8 appear multiplying off-diagonal Gell-Mann matrices in color space, meaning that they cannot be described as chemical potentials for suitable combinations of color charge densities. This in turn means that if any except μ_3 or μ_8 are nonzero, the analysis that we have set up in Section II is inapplicable. And, we have seen above that because flavor symmetry is explicitly broken in QCD, we *must* consider non-diagonal matrices Δ . It would seem that we are in trouble.

If we were trying to derive and solve gap equations for the patterns of pairing that we analyze, we would not

be able to evade the introduction of non-diagonal color “chemical potentials”. However, given our more limited goals, we can proceed as follows. Let Δ be an arbitrary gap matrix, in particular a non-diagonal one. For this gap matrix, the values of the color chemical potentials μ_a with $a = 1, \dots, 8$ required for color-neutrality can be determined explicitly as described in Appendix B. We set the μ_a ’s to these values, and now have a color-neutral state. We can now apply any color rotation, after which we will still have a color neutral state. So, we make that color rotation U that diagonalizes the matrix $\mu_a T_a$, where the T_a are the Gell-Mann matrices. Since $\mu_a T_a$ is Hermitian, such a transformation always exists. Moreover, since $\mu_a T_a$ is traceless, the diagonalized matrix is also traceless and thus can be written as a linear combination of T_3 and T_8 ,

$$U \mu_a T_a U^\dagger = \mu'_3 T_3 + \mu'_8 T_8, \quad (32)$$

where the new chemical potentials are functions of the old ones, $\mu'_3 = \mu'_3(\mu_1, \dots, \mu_8)$, $\mu'_8 = \mu'_8(\mu_1, \dots, \mu_8)$. The transformation U defines a transformation for all Gell-Mann matrices, $T_a \rightarrow U T_a U^\dagger$, as well as for the gap matrix, $\mathcal{M} \rightarrow (U \otimes \mathbf{1}) \mathcal{M}$ or, equivalently, $\Delta \rightarrow U^T \Delta$. Of course, all the new quantities depend on the chemical potentials $\mu_1, \dots, \mu_8$. As Eq. (B7) makes explicit, if the state was color neutral before this color rotation it is color neutral after it, assuming of course that the gap matrix is rotated by the same color rotation U as that applied to the chemical potentials. This means that for an arbitrary Δ we have found a physically equivalent order parameter $U^T \Delta$ for which color neutrality requires only nonzero μ_3 and μ_8 .

A more formal way of explaining what we have just done is to note that the Cartan subalgebra of $SU(3)_c$ has dimension 2, meaning that the maximal number of mutually commuting generators of $SU(3)_c$ is 2, and thus only two color charges can be measured simultaneously. This means that a phase with arbitrary pairing pattern Δ can be neutralized by the specification of only two linear combinations of the μ_a ’s. In general, these two combinations μ'_3 and μ'_8 are non-diagonal, involving all the μ_a ’s. However, by some color rotation U we find a new basis in which the two chemical potentials that must be nonzero are just μ_3 and μ_8 . Making this color rotation changes the pairing pattern from Δ to $U^T \Delta$. In general, neither Δ nor $U^T \Delta$ will be diagonal.

If we wanted to derive and solve gap equations, this result would be of little practical use as in order to find the “simpler” state $U^T \Delta$ explicitly one would have to have solved the whole problem for Δ . Our goals are more modest, however. We shall analyze the most general pairing pattern Δ , in the presence of only two color chemical potentials μ_3 and μ_8 , and show that stress-free pairing is impossible. If the Δ you find interesting is one of those which requires other μ_a ’s to be nonzero, we leave it to you to construct $U^T \Delta$. Since we will have analyzed all possible Δ ’s, your $U^T \Delta$ will be equal to another Δ that we have analyzed. Thus, our analysis below suffices.

V. STRESSED PAIRING IS UNAVOIDABLE

In this Section, we prove that for any possible pairing pattern in three-flavor quark matter, there will be a mismatch in the effective chemical potentials of at least two fermion species that form Cooper pairs with each other. We end the section with a discussion of some specific patterns of pairing in order to illustrate our method and the result by example.

A. General result

Here we shall apply the general discussion of Section II to the case of interest, namely nine fermions with effective chemical potentials as defined in Eq. (7) given by

$$\begin{aligned} \mu_{ru}^{\text{eff}} &= \mu - \frac{2}{3}\mu_e + \frac{1}{2}\mu_3 + \frac{1}{3}\mu_8, \\ \mu_{rd}^{\text{eff}} &= \mu + \frac{1}{3}\mu_e + \frac{1}{2}\mu_3 + \frac{1}{3}\mu_8, \\ \mu_{rs}^{\text{eff}} &= \mu + \frac{1}{3}\mu_e + \frac{1}{2}\mu_3 + \frac{1}{3}\mu_8 - \frac{m_s^2}{2\mu}, \\ \mu_{gu}^{\text{eff}} &= \mu - \frac{2}{3}\mu_e - \frac{1}{2}\mu_3 + \frac{1}{3}\mu_8, \\ \mu_{gd}^{\text{eff}} &= \mu + \frac{1}{3}\mu_e - \frac{1}{2}\mu_3 + \frac{1}{3}\mu_8, \\ \mu_{gs}^{\text{eff}} &= \mu + \frac{1}{3}\mu_e - \frac{1}{2}\mu_3 + \frac{1}{3}\mu_8 - \frac{m_s^2}{2\mu}, \\ \mu_{bu}^{\text{eff}} &= \mu - \frac{2}{3}\mu_e - \frac{2}{3}\mu_8, \\ \mu_{bd}^{\text{eff}} &= \mu + \frac{1}{3}\mu_e - \frac{2}{3}\mu_8, \\ \mu_{bs}^{\text{eff}} &= \mu + \frac{1}{3}\mu_e - \frac{2}{3}\mu_8 - \frac{m_s^2}{2\mu}. \end{aligned} \quad (33)$$

Electric neutrality is ensured via the electric chemical potential μ_e , while color neutrality is ensured via the color chemical potentials μ_3 and μ_8 , where we have normalized the corresponding Gell-Mann matrices as $T_3 = \text{diag}(1/2, -1/2, 0)$ and $T_8 = \text{diag}(1/3, 1/3, -2/3)$. Because $\text{Tr } Q = \text{Tr } T_3 = \text{Tr } T_8 = 0$, the condition (3) is satisfied. We consider color superconducting phases described by the gap matrix (29), specified by the 3×3 matrix Δ according to (28). We may now proceed along the same lines as for the toy models in Sec. III. The only difference is the larger number of fermion species and the resulting larger number of possible gap matrices. Because our gap matrix is specified by the nine parameters of Δ , and because we are only concerned about which parameters are zero and which nonzero, we have $2^9 = 512$ patterns of pairing to consider. In order to handle this large number of patterns, we make use of a one-to-one correspondence between a gap matrix (more precisely, between the pattern of zero and nonzero entries in a gap matrix) and a graph. We can then use *Maple* routines that know the basic terms of graph theory and which, in

conclusion.

B. Specific examples

We will illustrate our result using as examples gap matrices Δ for which we know that color neutrality can be imposed by adjusting only μ_3 and μ_8 , leaving $\mu_1 = \mu_2 = \mu_4 = \dots = \mu_7 = 0$. According to the results of Appendix B 2, we may consider all matrices which have at most one nonzero entry per column. By choosing the entries of Δ either zero or one, this gives rise to 64 gap matrices (each column may either be completely zero or may have a 1 at three possible entries, hence there are $4^3 = 64$ options). As we prove in Appendix C, 14 of these gap matrices are independent while all the others can be obtained from one of these 14 via a color rotation. In Table II we list these 14 matrices.

The left half of Table II contains all diagonal order parameters. One of them, here denoted by Δ_7 , is the pattern of pairing for the CFL phase. The corresponding graph is shown in the left panel of Fig. 4. The graphs of all the other diagonal order parameters in Table II are subgraphs of that for Δ_7 shown in Fig. 4, i.e., they are obtained by removing one or more edges of this graph. The left half of the table includes several phases that are familiar from the literature. For example, the Δ_3 pattern of pairing is the 2SC phase [33, 34], the uSC

and dSC phases described in Refs. [13, 46] are described by the patterns Δ_6 and Δ_5 , and the Δ_2 pattern is the 2SCus phase [6, 10]. The right half of Table II shows the seven patterns of pairing among our 14 examples which have non-diagonal gap matrices. If flavor symmetry were unbroken, each of these patterns would be equivalent to one of the diagonal ones. As flavor symmetry is explicitly broken, however, these patterns of pairing are physically distinct from the ones in the left half of the table. The electric and color chemical potentials in the table were obtained from the neutrality conditions (15).

Let us discuss two of the patterns of pairing in more detail, explicitly showing that the results for the chemical potentials in Table II do not allow for stress-free pairing. We do so for the patterns Δ_7 (CFL) and Δ_8 . The corresponding graphs are shown in Fig. 4. We take the explicit results from the table for μ_e , μ_3 , and μ_8 and use them to evaluate the effective chemical potentials of Eqs. (33). For the pattern of pairing Δ_8 , we find (showing only the components with more than a single quark)

$$\mathcal{C}_1 : \begin{aligned} \mu_{gu}^{\text{eff}} &= \mu - \frac{m_s^2}{6\mu} \\ \mu_{gd}^{\text{eff}} &= \mu + \frac{m_s^2}{12\mu} \\ \mu_{bs}^{\text{eff}} &= \mu - \frac{5m_s^2}{12\mu} \end{aligned}, \quad \mathcal{C}_2 : \begin{aligned} \mu_{gs}^{\text{eff}} &= \mu - \frac{5m_s^2}{12\mu} \\ \mu_{bu}^{\text{eff}} &= \mu - \frac{m_s^2}{6\mu} \\ \mu_{bd}^{\text{eff}} &= \mu + \frac{m_s^2}{12\mu} \end{aligned}. \quad (34)$$

It is obvious that this state does not exhibit stress-free pairing. For the CFL pattern Δ_7 , we obtain

$$\begin{aligned} \mathcal{C}_1 : \quad \mu_{ru}^{\text{eff}} &= \mu_{gd}^{\text{eff}} = \mu_{bs}^{\text{eff}} = \mu - \frac{m_s^2}{6\mu}, & \mathcal{C}_2 : \quad \mu_{rd}^{\text{eff}} &= \mu + \mu_e - \frac{m_s^2}{6\mu}, \\ & & \mu_{gu}^{\text{eff}} &= \mu - \mu_e - \frac{m_s^2}{6\mu}, \\ \mathcal{C}_3 : \quad \mu_{rs}^{\text{eff}} &= \mu - \frac{2m_s^2}{3\mu} + \mu_e, & \mathcal{C}_4 : \quad \mu_{gs}^{\text{eff}} &= \mu - \frac{2m_s^2}{3\mu}, \\ & & \mu_{bd}^{\text{eff}} &= \mu + \frac{m_s^2}{3\mu}. \end{aligned} \quad (35)$$

Even though one linear combination of μ_e , μ_3 and μ_8 is not fixed by the neutrality conditions, reflected here by the unspecified value of μ_e , we see that we can nevertheless draw a firm conclusion. We see that the fourth component, $\mathcal{C}_4$, exhibits a mismatch of effective chemical potentials for any value of μ_e , meaning that stress-free pairing is impossible. (In a treatment that goes beyond the order to which we are working, the electron chemical potential can be determined by including the electron pressure into the free energy. The result in the CFL case is $\mu_e = 0$ [6, 37], leading to the values for the effective chemical potentials illustrated in Fig. 1.) There are many patterns of pairing among the 511 that we have analyzed for which, at least at the order at which we are working, one combination of the chemical potentials is not fixed by the neutrality conditions. In all such cases we have shown that stress-free pairing cannot arise for any value

of the undetermined combination of chemical potentials.

VI. CONCLUSION

We have investigated all possible patterns of conventional BCS pairing in color-superconducting three-flavor quark matter in which the Cooper pairs are antisymmetric in color and flavor. We have analyzed 511 patterns of pairing, but it should be noted that many of them correspond to more than one phase, since our analysis only keeps track of which entries in the gap matrix are zero and which nonzero, not whether different nonzero entries are equal or unequal. Also, pushing in the opposite direction, not all of the 511 patterns are distinct: some are related by color rotation. Regardless, our analysis applies to every possible pattern, only a handful of which

diagonal order parameters	μ_e	μ_3	μ_8	non-diagonal order parameters	μ_e	μ_3	μ_8
$\Delta_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$	$\frac{m_s^2}{4\mu}$	0	0	$\Delta_8 = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$	$\frac{m_s^2}{4\mu}$	0	0
$\Delta_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$	0	0	0	$\Delta_9 = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$	$\frac{m_s^2}{4\mu}$	0	0
$\Delta_3 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	$\frac{m_s^2}{2\mu}$	0	0	$\Delta_{10} = \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$	$\frac{m_s^2}{4\mu}$	0	0
$\Delta_4 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$	0	0	$-\frac{m_s^2}{2\mu}$	$\Delta_{11} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$	μ_e	0	$-\mu_e + \frac{m_s^2}{\mu}$
$\Delta_5 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	$\frac{m_s^2}{2\mu}$	$\frac{m_s^2}{2\mu}$	$-\frac{m_s^2}{4\mu}$	$\Delta_{12} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$	μ_e	0	$-\mu_e - \frac{m_s^2}{2\mu}$
$\Delta_6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	μ_e	$\mu_e - \frac{m_s^2}{3\mu}$	$\frac{\mu_e}{2}$	$\Delta_{13} = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$	μ_e	0	$2\mu_e - \frac{m_s^2}{2\mu}$
$\Delta_7 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$	μ_e	μ_e	$\frac{\mu_e}{2} - \frac{m_s^2}{2\mu}$	$\Delta_{14} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$	$\frac{m_s^2}{4\mu}$	$-\frac{2}{3}\mu_8$	μ_8

TABLE II: Order parameters for which $\mu_1 = \mu_2 = \mu_4 = \dots = \mu_7 = 0$. The electric and color chemical potentials satisfying the neutrality constraints are given for each pattern of pairing. In those cases in which the neutrality conditions leave one linear combination of μ_e , μ_3 and μ_8 undetermined, we write the chemical potentials in terms of an unspecified μ_e (or μ_8 in the one case in which the undetermined combination does not involve μ_e .)

have been studied before.

At asymptotic densities, where the differences among the quark masses can be neglected, previous work has shown that the CFL phase is favored [1]. As the density is decreased, CFL pairing requires that the system pay a free energy cost that is parametrically of order $(m_s^2/\mu)^2\mu^2 = m_s^4$, in order to gain a free energy benefit of order $\Delta^2\mu^2$. This guarantees that below some μ of

order m_s^2/Δ , the CFL phase must break down. In this investigation we have shown that *all* 511 possible pairing patterns suffer the same fate. All of them exact a free energy cost of order m_s^4 ; none are stress-free.

There are, then, three classes of possible answers to the question of what succeeds the CFL phase as a function of decreasing density. First, if m_s^2/Δ is small enough the CFL phase will be succeeded directly by nuclear matter:




FIG. 4: Graphs for the patterns of pairing Δ_7 (CFL) and Δ_8 , cf. Table II.

the quark matter $\rightarrow$ nuclear matter transition will occur before μ drops to the critical value of order m_s^2/Δ at which CFL pairing breaks down. The second possibility is that the CFL phase is succeeded by a color-superconducting state with unconventional pairing, not simply as in BCS theory. Such a state could be one which is spatially inhomogeneous, one which includes quasiparticles which are paired but gapless, one which includes counterpropagating currents, one which includes meson condensates, or one which involves any combination of these possibilities. Third, it is a logical possibility that the CFL phase is succeeded by some among the 510 less symmetrically paired BCS phases that we have considered, for example the 2SC phase. However, the conclusion of our work is that any of these must also break down below a μ of order m_s^2/Δ , meaning that any of these must also be succeeded by some state with non-BCS pairing. None of the 511 possible patterns of BCS pairing can survive in a regime in which $\mu \ll m_s^2/\Delta$.

Acknowledgments

We thank E. Gubankova, H. Malekzadeh, M. Mannarelli, D. Rischke, R. Sharma, and I. Shovkovy for valuable comments and discussions. AS acknowledges financial support from the German Academic Exchange Service (DAAD). KR acknowledges the hospitality of the Nuclear Theory Group at LBNL. This research was supported in part by the Director, Office of Science, Office of Nuclear Physics, of the U.S. Department of Energy under contract #DE-AC02-05CH11231 and cooperative

research agreement #DF-FC02-94ER40818.

APPENDIX A: WHY THERE IS NO “A PHASE” IN THE CASE OF A $SU(3)_c \times SU(3)_f$ SYMMETRY

In this Appendix, we elaborate on the difference between a three-flavor color superconductor on the one hand and superfluid ^3He and a one-flavor color superconductor on the other hand. All three systems are described by an order parameter parametrized by a 3×3 matrix Δ . The reason is that in all cases the Cooper pairing occurs in channels which are triplets with respect to two different symmetries. In a three-flavor color superconductor these are the color and flavor (anti)triplets, in superfluid ^3He the angular momentum and spin triplets, and in a one-flavor color superconductor the color and total angular momentum (anti)triplets. In the latter two systems there is a non-diagonal order parameter matrix

$$\Delta_A = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & i & 0 \end{pmatrix}, \quad (\text{A1})$$

describing the so-called *A* phase [43, 44]. This phase is genuinely different from all phases that can be described by a diagonal matrix Δ because there is no symmetry rotation that transforms Δ_A into a diagonal matrix. In the case of ^3He , the symmetry rotations are elements of the group $SO(3)_L \times SO(3)_S$, where $SO(3)_L$ and $SO(3)_S$ are the angular momentum and spin groups, respectively. This group acts on Δ as given in Eq. (31). Suppose

there is a transformation $U \otimes V$ with $U \in SO(3)_L$ and $V \in SO(3)_S$ that diagonalizes Δ_A . This requires that the off-diagonal elements of the resulting matrix vanish,

$$\sum_{k\ell} u_{ki} \Delta_{k\ell} v_{\ell j} = 0 \quad \text{for } i \neq j, \quad (\text{A2})$$

where u_{ki} and $v_{\ell j}$ are the entries of the matrices U and V respectively. Or, explicitly,

$$u_{31}(v_{12} + iv_{22}) = u_{31}(v_{13} + iv_{23}) = 0, \quad (\text{A3a})$$

$$u_{32}(v_{11} + iv_{21}) = u_{32}(v_{13} + iv_{23}) = 0, \quad (\text{A3b})$$

$$u_{33}(v_{11} + iv_{21}) = u_{33}(v_{12} + iv_{22}) = 0. \quad (\text{A3c})$$

Since U is unitary and thus invertible, at least one of the entries u_{31} , u_{32} , and u_{33} must be nonzero. Without loss of generality, we may assume that $u_{31} \neq 0$. Since all v_{ij} 's are real, this means that $v_{12} = v_{22} = v_{13} = v_{23} = 0$. Therefore,

$$V V^T = \begin{pmatrix} v_{11}^2 & v_{21}v_{11} & v_{31}v_{11} \\ v_{21}v_{11} & v_{21}^2 & v_{31}v_{21} \\ v_{31}v_{11} & v_{31}v_{21} & v_{31}^2 + v_{32}^2 + v_{33}^2 \end{pmatrix}. \quad (\text{A4})$$

The orthogonality of V then implies $v_{11}^2 = v_{21}^2 = 1$ and $v_{21}v_{11} = 0$. These two conditions contradict each other. Consequently, there is no transformation $U \otimes V$ that diagonalizes Δ_A .

Since we only needed the v_{ij} 's to be real, the same proof holds for complex u_{ij} 's, i.e., for the symmetry group $SU(3)_c \times SO(3)_J$, relevant for a one-flavor color superconductor, where J is the total angular momentum.

However, in the case of the symmetry group $SU(3)_c \times SU(3)_f$, hence for unitary transformations U and V , any matrix Δ can be diagonalized by $U^T \Delta V$, see for example Ref. [42]. In particular, a potential “ A phase” order parameter is diagonalized by the pure flavor rotation

$$U = \mathbf{1}, \quad V = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & i & 1 \\ 0 & -1 & -i \\ \sqrt{2} & 0 & 0 \end{pmatrix}, \quad (\text{A5})$$

where $U \in SU(3)_c$ and $V \in SU(3)_f$. With these matrices we find

$$U^T \Delta_A V = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (\text{A6})$$

Hence, if $SU(3)_c \times SU(3)_f$ were unbroken in three-flavor QCD, the “ A phase” would be equivalent to one with a diagonal Δ .

As described in Section IV and manifest in Section V, however, the explicit breaking of $SU(3)_f$ in QCD by a nonzero strange quark mass means that a generic non-diagonal Δ is *not* equivalent to a diagonal Δ . We must therefore consider all 511 patterns of pairing including those which are analogues of the A phase.

APPENDIX B: TADPOLES AND COLOR NEUTRALITY

In this Appendix we determine which color chemical potentials have to be nonzero to neutralize a three-flavor color superconducting phase whose gap matrix $\mathcal{M}$ is specified by (28), with Δ an arbitrary 3×3 matrix. In QCD, these chemical potentials originate from nonzero expectation values of the corresponding zeroth component of the gluon fields A_a^0 , $a = 1, \dots, 8$. In the mean-field approximation, the Yang-Mills equation for the gluon fields requires the zeroth component of the quark tadpole contributions to vanish [38, 39],

$$\mathcal{T}_a^0 = 0, \quad a = 1, \dots, 8. \quad (\text{B1})$$

Here, “quark tadpole” refers to the Feynman diagram of a quark loop with a gluon leg. The explicit expressions are given by [39]

$$\mathcal{T}_a^\mu = -\frac{g}{2} \text{Tr}[\Gamma_a^\mu \mathcal{S}], \quad (\text{B2})$$

where g is the strong coupling constant. Before discussing the explicit structure of the quark propagator $\mathcal{S}$, let us point out the physical meaning of the eight equations (B1). The two tadpoles $\mathcal{T}_3$ and $\mathcal{T}_8$ are proportional to color density differences, and even though the other tadpoles cannot be interpreted in this way this makes clear that satisfying the requirement that all tadpoles vanish results in a color-neutral state. One often replaces the precise statement that the eight μ_a 's take on values such that the eight $\mathcal{T}_a$'s vanish by the loose statement that color neutrality is achieved by adjusting the color chemical potentials μ_a such that all color densities vanish. As explained in Section IV, this loose statement can always be rendered precise by a color rotation, which also rotates Δ .

In normal quark matter, the tadpoles vanish even with vanishing gluon fields $A_1^0 = \dots = A_8^0 = 0$. In superconducting quark matter, however, nonvanishing gluon fields have to be introduced in order to fulfil Eqs. (B1). Which gluon fields attain a nonzero expectation value depends on the pairing pattern.

1. Formalism

In Eq. (B2), the vertex $\Gamma_a^\mu \equiv \text{diag}(\gamma^\mu T_a, -\gamma^\mu T_a^T)$, where T_a are the Gell-Mann matrices, and the quark propagator

$$\mathcal{S} = \begin{pmatrix} G^+ & \Xi^- \\ \Xi^+ & G^- \end{pmatrix} \quad (\text{B3})$$

are given in the Nambu-Gorkov basis. We do not need to specify the exact form of the anomalous propagators $\Xi^\pm$ because they drop out in the expression for the tadpole after taking trace over Nambu-Gorkov space. The

diagonal components $G^\pm$ are the propagators for quarks and charge-conjugated quarks,

$$G^\pm \equiv \{[G_0^\pm]^{-1} - \Phi^\mp G_0^\mp \Phi^\pm\}^{-1}, \quad (\text{B4})$$

where $G_0^\pm$ is the tree-level quark (charge-conjugate quark) propagator and $\Phi^\mp$ the off-diagonal components of the quark self-energy. In general, in order to ensure color neutrality, all 8 color chemical potentials enter the (inverse) tree-level propagators,

$$[G_0^\pm]^{-1} = \gamma^\mu K_\mu \pm \gamma_0 (\mu + \mu_a T_a + \mu_e Q) - m, \quad (\text{B5})$$

where summation over $a = 1, \dots, 8$ is implied, and K_μ is the quark four-momentum. The matrices Q and m are proportional to the identity in color space, and are diagonal in flavor space. They account for the quark electric charges and masses, respectively. For the following qualitative discussion, we are not interested in the exact value of the tadpoles. We are rather interested in the effect of the color-flavor structure on $\mathcal{T}_a^0$. Thus, we may drop all Dirac and Nambu-Gorkov structure and write the tadpoles as

$$\mathcal{T}_a^0 = \text{Tr} \left\{ T_a [F + \mu_b T_b - \mathcal{M}^\dagger (F - \mu_b T_b)^{-1} \mathcal{M}]^{-1} \right\}, \quad (\text{B6})$$

where F is a diagonal matrix in flavor space. We have replaced the (inverse) propagator by the schematic expression $[G_0^\pm]^{-1} \rightarrow F \pm \mu_a T_a$ and $\Phi^+ \rightarrow \mathcal{M}$, $\Phi^- \rightarrow \mathcal{M}^\dagger$.

We remark that the tadpoles remain invariant under color rotations of the Gell-Mann matrices and the gap matrix,

$$\mathcal{T}_a^0 [T_a, \mathcal{M}] = \mathcal{T}_a^0 [U T_a U^\dagger, U \mathcal{M}]. \quad (\text{B7})$$

Although this is not difficult to see, we present here an explicit proof. The proof is basically a repeated application of the formula

$$(F + U M U^\dagger)^{-1} = U (F + M)^{-1} U^\dagger, \quad (\text{B8})$$

which is valid for unitary U and $U F U^\dagger = F$. In the following, F is a matrix in flavor space, i.e., $F = \mathbf{1} \otimes F$, and U acts on color space, i.e., $U F U^\dagger = U U^\dagger \otimes F = F$; M can be a matrix in both color and flavor spaces, say $M = m_{ij} A_i \otimes B_j$ with complex coefficients m_{ij} and color and flavor matrices A_i and B_j , respectively. Then, by making use of Eq. (30),

$$\begin{aligned} \mathcal{T}_a^0 [U T_a U^\dagger, U \mathcal{M}] &= \text{Tr} \left\{ U T_a U^\dagger [F + \mu_b U T_b U^\dagger - \Delta_{ij}^* \Delta_{mn} U J_i U^\dagger I_j (F - \mu_b U T_b U^\dagger)^{-1} U J_m U^\dagger I_n]^{-1} \right\} \\ &= \text{Tr} \left\{ U T_a U^\dagger [F + \mu_b U T_b U^\dagger - \Delta_{ij}^* \Delta_{mn} U J_i I_j (F - \mu_b T_b)^{-1} J_m U^\dagger I_n]^{-1} \right\} \\ &= \text{Tr} \left\{ U T_a U^\dagger [F + U \{ \mu_b T_b - \mathcal{M}^\dagger (F - \mu_b T_b)^{-1} \mathcal{M} \} U^\dagger]^{-1} \right\} \\ &= \text{Tr} \left\{ T_a [F + \mu_b T_b - \mathcal{M}^\dagger (F - \mu_b T_b)^{-1} \mathcal{M}]^{-1} \right\} \\ &= \mathcal{T}_a^0 [T_a, \mathcal{M}], \end{aligned} \quad (\text{B9})$$

which is Eq. (B7).

2. Evaluating tadpoles

We may now use the general form of the tadpoles (B6) to investigate which chemical potentials are needed for a given gap matrix Δ . To this end, we have to evaluate the tadpoles for *vanishing* color chemical potentials $\mu_1 = \dots = \mu_8 = 0$. Let us denote these quantities by $\mathcal{T}_a^0(0)$. From Eq. (B6) we obtain

$$\mathcal{T}_a^0(0) \equiv \text{Tr} \left[T_a (F - \mathcal{M}^\dagger F^{-1} \mathcal{M})^{-1} \right], \quad (\text{B10})$$

where $F \equiv \text{diag}(f_1, f_2, f_3)$ accounts for the explicit flavor symmetry breaking. A nonvanishing tadpole, $\mathcal{T}_a^0(0) \neq 0$, indicates that the corresponding color chemical potential

μ_a has to be nonzero in order to obtain a color neutral state. The converse, namely the statement that a vanishing tadpole, $\mathcal{T}_a^0(0) = 0$, indicates that the state can be color-neutralized with the corresponding color chemical potential μ_a set to zero, is not immediate. We have not found a general argument that precludes the concern that after introducing a nonzero color chemical μ_a that cancels the initially nonzero $\mathcal{T}_a^0(0)$, some other tadpole $\mathcal{T}_b^0$ could now be nonvanishing, even though $\mathcal{T}_b^0(0)$ was zero. We have, however, checked that for very many pairing patterns, including all those in Table II and many more, this concern does not arise. We have checked this for patterns with multiple nonzero entries per column in Δ , with complex entries in Δ , and for patterns with up to five different nonzero $\mathcal{T}_a^0(0)$'s, which of course means nonzero $\mathcal{T}_a^0(0)$'s that do not commute. This suggests the following conjecture: First, use Eq. (B10) to compute the

tadpoles with all the μ_a 's set to zero, namely the $\mathcal{T}_a^0(0)$ for all $a = 1, \dots, 8$. Now, keep $\mu_a = 0$ only for those values of a for which $\mathcal{T}_a^0(0) = 0$; for the other values of a , reintroduce μ_a as a free variable. Then, use Eq. (B6) to compute all eight tadpoles $\mathcal{T}_b^0$ as a function of the nonzero μ_a 's. We conjecture that, for all pairing patterns Δ , the $\mathcal{T}_b^0$ computed in this way will vanish for all values of b for which $\mathcal{T}_b^0(0)$ was found to vanish in the first step. We do not have a proof of this conjecture, but we know of no counterexample and, as described above, we have checked many examples.

Having described the consequences of which $\mathcal{T}_a^0(0)$ are zero and which nonzero, we turn now to the evaluation of the expression (B10). We begin with the simplest possible situation, setting $F = \mathbf{1}$, i.e. switching off all flavor symmetry breaking terms. We are left with the inversion of the matrix $\mathbf{1} - \mathcal{M}^\dagger \mathcal{M}$. It can be formally inverted by writing the Hermitian matrix $\mathcal{M}^\dagger \mathcal{M}$ in terms of its eigenvalues λ_i and the projectors onto the corresponding eigenspaces $\mathcal{P}_i$,

$$\mathcal{M}^\dagger \mathcal{M} = \sum_{i=1}^n \lambda_i \mathcal{P}_i, \quad (\text{B11})$$

where we denoted the number of different eigenvalues by n . Then, since $\{\mathcal{P}_1, \dots, \mathcal{P}_n\}$ is a complete set of orthogonal projectors,

$$(\mathbf{1} - \mathcal{M}^\dagger \mathcal{M})^{-1} = \sum_{i=1}^n \frac{1}{1 - \lambda_i} \mathcal{P}_i. \quad (\text{B12})$$

The projectors are given by (see for example Appendix A of Ref. [47])

$$\mathcal{P}_i = \prod_{j \neq i}^n \frac{\mathcal{M}^\dagger \mathcal{M} - \lambda_j}{\lambda_i - \lambda_j}. \quad (\text{B13})$$

As a further simplification, let us assume that $\mathcal{M}^\dagger \mathcal{M}$ has at most two different eigenvalues. Then, the projectors $\mathcal{P}_i$ are a linear combination of the matrices $\mathbf{1}$ and $\mathcal{M}^\dagger \mathcal{M}$ (in the case of more than two different eigenvalues, higher powers of $\mathcal{M}^\dagger \mathcal{M}$ are present). Since $\text{Tr } T_a = 0$, we obtain

$$\mathcal{T}_a^0(0) \propto \text{Tr}[T_a \mathcal{M}^\dagger \mathcal{M}] = -2 \text{Tr}[T_a \Delta \Delta^\dagger], \quad (\text{B14})$$

where Eq. (28) as well as $\text{Tr}[T_a J_i J_j] = -(T_a)_{ij}$ and $\text{Tr}[I_i I_j] = 2\delta_{ij}$ have been used. This simple form of the tadpole can be applied for example to the CFL phase with $m_s = 0$. In this case $\Delta = \mathbf{1}$ and the eigenvalues of $\mathcal{M}^\dagger \mathcal{M}$ are 4 and 1. We immediately conclude $\mathcal{T}_a^0(0) = 0$ for all a . This reflects the fact that in the absence of flavor symmetry breaking the CFL phase is automatically color neutral, i.e., there is no nonzero color chemical potential.

The simplest way to introduce explicit flavor symmetry breaking “by hand” is to insert a flavor matrix F into Eq. (B14),

$$\mathcal{T}_a^0(0) \propto \text{Tr}[T_a \mathcal{M}^\dagger F \mathcal{M}] = -\text{Tr}[T_a \Delta F' \Delta^\dagger], \quad (\text{B15})$$

where we used $\text{Tr}[I_i F I_j] = \delta_{ij} \text{Tr } F - F_{ij}$ and defined $F' \equiv \text{Tr}[F] \cdot \mathbf{1} - F$. (Since the only purpose of F' is to break the flavor symmetry, we may drop the prime in the following.) The expressions for the tadpoles can now be evaluated explicitly:

$$\mathcal{T}_1^0(0) \propto f_1 \text{Re}[\Delta_{11} \Delta_{21}^*] + f_2 \text{Re}[\Delta_{12} \Delta_{22}^*] + f_3 \text{Re}[\Delta_{13} \Delta_{23}^*] \quad (\text{B16a})$$

$$\mathcal{T}_2^0(0) \propto i(f_1 \text{Im}[\Delta_{11} \Delta_{21}^*] + f_2 \text{Im}[\Delta_{12} \Delta_{22}^*] + f_3 \text{Im}[\Delta_{13} \Delta_{23}^*]) \quad (\text{B16b})$$

$$\mathcal{T}_3^0(0) \propto \frac{1}{2}[f_1(|\Delta_{11}|^2 - |\Delta_{21}|^2) + f_2(|\Delta_{12}|^2 - |\Delta_{22}|^2) + f_3(|\Delta_{13}|^2 - |\Delta_{23}|^2)] \quad (\text{B16c})$$

$$\mathcal{T}_4^0(0) \propto f_1 \text{Re}[\Delta_{11} \Delta_{31}^*] + f_2 \text{Re}[\Delta_{12} \Delta_{32}^*] + f_3 \text{Re}[\Delta_{13} \Delta_{33}^*] \quad (\text{B16d})$$

$$\mathcal{T}_5^0(0) \propto i(f_1 \text{Im}[\Delta_{11} \Delta_{31}^*] + f_2 \text{Im}[\Delta_{12} \Delta_{32}^*] + f_3 \text{Im}[\Delta_{13} \Delta_{33}^*]) \quad (\text{B16e})$$

$$\mathcal{T}_6^0(0) \propto f_1 \text{Re}[\Delta_{21} \Delta_{31}^*] + f_2 \text{Re}[\Delta_{22} \Delta_{32}^*] + f_3 \text{Re}[\Delta_{23} \Delta_{33}^*] \quad (\text{B16f})$$

$$\mathcal{T}_7^0(0) \propto i(f_1 \text{Im}[\Delta_{21} \Delta_{31}^*] + f_2 \text{Im}[\Delta_{22} \Delta_{32}^*] + f_3 \text{Im}[\Delta_{23} \Delta_{33}^*]) \quad (\text{B16g})$$

$$\begin{aligned} \mathcal{T}_8^0(0) \propto & \frac{1}{2\sqrt{3}}[f_1(|\Delta_{11}|^2 + |\Delta_{21}|^2 - 2|\Delta_{31}|^2) + f_2(|\Delta_{12}|^2 + |\Delta_{22}|^2 - 2|\Delta_{32}|^2) \\ & + f_3(|\Delta_{13}|^2 + |\Delta_{23}|^2 - 2|\Delta_{33}|^2)] . \end{aligned} \quad (\text{B16h})$$

Note that we have obtained Eqs. (B16) by inserting flavor-breaking by hand, and have not been able to de-

rive them from the rigorous expression (B10). However, by inserting various gap matrices (including all the ex-

amples in Table II and many more) explicitly into Eq. (B10), we have checked that (B16) and the rigorous form (B10) yield the same conclusion. I.e., a given gap matrix gives rise to the same nonzero tadpoles $\mathcal{T}_a^0(0)$ whether we use (B10) or (B16). We can also use the two-flavor result given below in Eqs. (B17) as another check: Eqs. (B17) can be obtained from Eqs. (B16) immediately, and we *have* been able to derive (B17) rigorously.

As an example of the use of (B16), we can now apply these equations to the CFL and gCFL phases, in both of which the only nonzero Δ 's are Δ_{11} , Δ_{22} and Δ_{33} . From this we see immediately that in the CFL and gCFL phases, the tadpoles $\mathcal{T}_1^0(0)$, $\mathcal{T}_2^0(0)$, $\mathcal{T}_4^0(0)$, $\dots$, $\mathcal{T}_7^0(0)$ vanish whereas the tadpoles $\mathcal{T}_3^0(0)$ and $\mathcal{T}_8^0(0)$ may be nonzero due to the explicit flavor symmetry breaking. In fact, in the CFL phase $f_1 = f_2$ and $|\Delta_{11}| = |\Delta_{22}|$, which makes $\mathcal{T}_3^0(0) = 0$ leaving only $\mathcal{T}_8^0(0)$ nonzero. This is consistent with the result of Refs. [6, 7] that the CFL phase is color neutral with $\mu_3 = 0$ and $\mu_8 = -m_s^2/(2\mu)$. In the gCFL phase, however, $|\Delta_{11}| \neq |\Delta_{22}|$ and both $\mathcal{T}_3^0(0)$ and $\mathcal{T}_8^0(0)$ are nonzero. Consequently, Eqs. (B16) indicate that in the gapless CFL phase color neutrality can be ensured via the introduction of suitable nonzero values of μ_3 and μ_8 , as was shown explicitly in Refs. [9, 10]. Furthermore, Eqs. (B16) show that $\mu_1, \mu_2, \mu_4 \dots \mu_7$ vanish, as assumed in Refs. [9, 10, 11, 12, 13, 14, 15, 16] and demonstrated in Ref. [48].

We can apply Eqs. (B16) more generally as follows. We note that for (a, b) being either $(1, 2)$, $(4, 5)$, or $(6, 7)$ the following statement holds: both $\mathcal{T}_a$ and $\mathcal{T}_b$ are zero if at least one member of each of the following three pairs is zero: $(\Delta_{m1}, \Delta_{n1})$, $(\Delta_{m2}, \Delta_{n2})$, $(\Delta_{m3}, \Delta_{n3})$. Here, $(m, n) = (1, 2)$ for $(a, b) = (1, 2)$, $(m, n) = (1, 3)$ for $(a, b) = (4, 5)$, and $(m, n) = (2, 3)$ for $(a, b) = (6, 7)$. (The argument is straightforward. Consider one of these pairs, say $(\Delta_{m1}, \Delta_{n1})$. To make both $\mathcal{T}_a^0(0)$ and $\mathcal{T}_b^0(0)$ vanish, according to Eqs. (B16) we have to require $\text{Re}[\Delta_{m1}\Delta_{n1}^*] = \text{Im}[\Delta_{m1}\Delta_{n1}^*] = 0$. Hence $\Delta_{m1}\Delta_{n1}^*$ and therefore either Δ_{m1} or Δ_{n1} (or both) must vanish.) Taking the case $(a, b) = (1, 2)$ as an example, we have learned that both $\mathcal{T}_1^0(0)$ and $\mathcal{T}_2^0(0)$ vanish if either the first or the second (or both) entries in each column of the order parameter matrix Δ vanish. Including the other two cases, we reach the following conclusion: $\mathcal{T}_1^0(0) = \mathcal{T}_2^0(0) = \mathcal{T}_4^0(0) = \dots = \mathcal{T}_7^0(0) = 0$ if each column of the matrix Δ contains at most one nonzero entry.

We can also derive an analogous result for a two-flavor color superconductor. This case differs from the current one in the structure of the order parameter. Since condensation takes place in the flavor-singlet channel, the order parameter is given by a three-vector $\Delta \equiv (\Delta_1, \Delta_2, \Delta_3)$ than by a 3×3 matrix. The gap matrix is $\mathcal{M} = \Delta_i J_i \tau_2$, with the antisymmetric flavor singlet described by the second Pauli matrix τ_2 . Repeating the same steps as for the three-flavor case, we find $\text{Tr}[T_a \mathcal{M}^\dagger F \mathcal{M}] = -\text{Tr}[F] \Delta^\dagger T_a \Delta$, with $F \equiv \text{diag}(f_1, f_2)$,

and thus

$$\mathcal{T}_1^0(0) \propto (f_1 + f_2) \text{Re}[\Delta_1 \Delta_2^*] \quad (\text{B17a})$$

$$\mathcal{T}_2^0(0) \propto i(f_1 + f_2) \text{Im}[\Delta_1 \Delta_2^*] \quad (\text{B17b})$$

$$\mathcal{T}_3^0(0) \propto \frac{1}{2}(f_1 + f_2)(|\Delta_1|^2 - |\Delta_2|^2) \quad (\text{B17c})$$

$$\mathcal{T}_4^0(0) \propto (f_1 + f_2) \text{Re}[\Delta_1 \Delta_3^*] \quad (\text{B17d})$$

$$\mathcal{T}_5^0(0) \propto i(f_1 + f_2) \text{Im}[\Delta_1 \Delta_3^*] \quad (\text{B17e})$$

$$\mathcal{T}_6^0(0) \propto (f_1 + f_2) \text{Re}[\Delta_2 \Delta_3^*] \quad (\text{B17f})$$

$$\mathcal{T}_7^0(0) \propto i(f_1 + f_2) \text{Im}[\Delta_2 \Delta_3^*] \quad (\text{B17g})$$

$$\mathcal{T}_8^0(0) \propto \frac{f_1 + f_2}{2\sqrt{3}}(|\Delta_1|^2 + |\Delta_2|^2 - 2|\Delta_3|^2) \quad (\text{B17h})$$

We see that $\mathcal{T}_1^0(0) = \mathcal{T}_2^0(0) = \mathcal{T}_4^0(0) = \dots = \mathcal{T}_7^0(0) = 0$ if the vector Δ contains at most one nonzero entry.

The main results of this appendix can be summarized as follows:

- The tadpoles $\mathcal{T}_1^0(0)$, $\mathcal{T}_2^0(0)$, $\mathcal{T}_4^0(0)$, $\mathcal{T}_5^0(0)$, $\mathcal{T}_6^0(0)$, $\mathcal{T}_7^0(0)$ vanish if each column in the gap matrix Δ contains at most one nonzero entry. (In a two-flavor color superconductor, they vanish if the vector Δ contains at most one nonzero entry.)
- If there is at least one column in the gap matrix Δ with more than one nonzero entry, the “usual” color chemical potentials μ_3 and μ_8 are not sufficient to ensure color neutrality. (In the two flavor case, this arises if the vector Δ has more than one nonzero entry. The possibility of chemical potentials other than μ_3 and μ_8 in a two-flavor color superconductor has been discussed in detail in Ref. [45].)

Note that in the two flavor case, a color rotation can always be found that takes a generic gap vector Δ into the 2SC pattern $\Delta = (0, 0, 1)$, meaning that chemical potentials other than μ_3 and μ_8 can always be avoided. In the three flavor case, on the other hand, a color rotation alone cannot turn a generic gap matrix Δ into one which has only one nonzero entry per column. We must therefore proceed as described in Sections IV and V, making a color rotation that diagonalizes $\mu_a T_a$ but does not simplify Δ , meaning that we have to analyze an exhaustive list of Δ 's.

APPENDIX C: SUPERCONDUCTING PHASES IN TABLE II

In this Appendix, we prove that

1. Any of the 64 patterns of pairing described by 3×3 matrices Δ which have at most one nonzero entry per column can be related to one of the 14 patterns of pairing in Table II by a color rotation.
2. The 14 patterns of pairing in Table II are not connected by any color rotation.

To prove the first point, we have to show that we obtain all 64 patterns of pairing that have at most one nonzero entry per column by color rotations from the phases in Table II. One can simply apply a color permutation to each of the phases. A color permutation is equivalent to a permutation of the rows in the matrix Δ . Therefore, Δ_1 in Table II represents 3 matrices, and so do Δ_2 and Δ_3 . Δ_4 through Δ_7 represent 6 matrices each, etc. In total, in the same order as in Table II, $3+3+3+6+6+6+6+3+3+3+6+6+6+3=63$. The 64th is the zero matrix.

To prove the second point, we have to show that for any pair (Δ_i, Δ_j) chosen from among the 14 patterns of pairing in Table II, there is no unitary matrix $U \in SU(3)$ such that $\Delta_j = U \Delta_i$. This is obvious if there is one column that is completely zero in Δ_i but has a nonzero entry in Δ_j . In this case, there is no matrix — whether unitary or not — that can transform Δ_i into Δ_j . Therefore, the 14 matrices can be divided into the following subsets: $\{\Delta_1\}$, $\{\Delta_2\}$, $\{\Delta_3\}$, $\{\Delta_4, \Delta_8\}$, $\{\Delta_5, \Delta_9\}$, $\{\Delta_6, \Delta_{10}\}$, and $\{\Delta_7, \Delta_{11}, \Delta_{12}, \Delta_{13}, \Delta_{14}\}$. For any pair (Δ_i, Δ_j) with Δ_i and Δ_j chosen from two different subsets, the statement is obviously true. The remaining cases must be treated separately. Suppose there is a U such that $U \Delta_4 = \Delta_8$. It follows that $u_{21} = u_{22} = u_{31} = u_{32} = 0$, where u_{ij} are the entries of the matrix U . Then,

$$UU^\dagger = \begin{pmatrix} |u_{11}|^2 + |u_{12}|^2 + |u_{13}|^2 & u_{13}u_{23}^* & u_{13}u_{33}^* \\ u_{23}u_{13}^* & |u_{23}|^2 & u_{23}u_{33}^* \\ u_{33}u_{13}^* & u_{33}u_{23}^* & |u_{33}|^2 \end{pmatrix}. \quad (C1)$$

The requirement that U be unitary leads to the contradictory requirements $|u_{23}|^2 = |u_{33}|^2 = 1$ and $u_{23}u_{33}^* = 0$. Consequently, Δ_4 and Δ_8 are not equivalent. By symmetry, the same argument applies to the pairs (Δ_5, Δ_9) and (Δ_6, Δ_{10}) . Only the last of the subsets remains. Suppose that there is a U with $U \Delta_7 = \Delta_{11}$. This leads to a vanishing row in U , $u_{31} = u_{32} = u_{33} = 0$. Thus, U cannot be invertible and thus not unitary. With the same argument one shows that Δ_7 is not equivalent to Δ_{12} , Δ_{13} , and Δ_{14} . Now suppose that $U \Delta_{11} = \Delta_{12}$. Then, there must be a vanishing column in U , $u_{11} = u_{21} = u_{31} = 0$. Hence this U cannot be unitary. By the same argument, the equivalence of Δ_{11} , Δ_{13} and Δ_{12} , Δ_{13} is excluded. For $U \Delta_{11} = \Delta_{14}$ we find $u_{21} = u_{22} = u_{31} = u_{32} = 0$. As shown above, this cannot be fulfilled by a unitary matrix. Therefore, Δ_{11} and Δ_{14} are not equivalent, and, by symmetry, the same holds for the pairs $(\Delta_{12}, \Delta_{14})$ and $(\Delta_{13}, \Delta_{14})$. This proves that all 14 matrices are independent.

-
- [1] For reviews, see K. Rajagopal and F. Wilczek, arXiv:hep-ph/0011333; M. G. Alford, Ann. Rev. Nucl. Part. Sci. **51**, 131 (2001) [arXiv:hep-ph/0102047]; G. Nardulli, Riv. Nuovo Cim. **25N3**, 1 (2002) [arXiv:hep-ph/0202037]; S. Reddy, Acta Phys. Polon. B **33**, 4101 (2002) [arXiv:nucl-th/0211045]; T. Schäfer, arXiv:hep-ph/0304281; D. H. Rischke, Prog. Part. Nucl. Phys. **52**, 197 (2004) [arXiv:nucl-th/0305030]; M. Alford, Prog. Theor. Phys. Suppl. **153**, 1 (2004) [arXiv:nucl-th/0312007]; M. Buballa, Phys. Rept. **407**, 205 (2005) [arXiv:hep-ph/0402234]; H. c. Ren, arXiv:hep-ph/0404074; I. Shovkovy, arXiv:nucl-th/0410091; T. Schäfer, arXiv:hep-ph/0509068.
- [2] J. Bardeen, L.N. Cooper, and J.R. Schrieffer, Phys. Rev. **108**, 1175 (1957).
- [3] M. G. Alford, K. Rajagopal and F. Wilczek, Nucl. Phys. B **537**, 443 (1999) [arXiv:hep-ph/9804403].
- [4] K. Iida and G. Baym, Phys. Rev. D **63**, 074018 (2001) [Erratum-ibid. D **66**, 059903 (2002)] [arXiv:hep-ph/0011229];
- [5] Stable bulk matter must be neutral under all gauged charges, whether they are spontaneously broken or not. Otherwise, the net charge density would create large electric fields, making the energy non-extensive. In the case of the electromagnetic gauge symmetry, this simply requires zero charge density. In the case of the color gauge symmetry, bulk matter must in fact be a color singlet, which is a more restrictive condition than mere color neutrality. However, the free energy cost of projecting a color neutral state onto a color singlet state falls rapidly with volume, as long as we are considering volumes larger than the size of a Cooper pair. Given that if quark matter occurs within the core of a neutron star the relevant volumes will be of order cubic kilometers, whereas Cooper pairs have sizes of order fm, it is more than sufficient to consider only the consequences of enforcing color neutrality. See P. Amore, M. C. Birse, J. A. McGovern and N. R. Walet, Phys. Rev. D **65**, 074005 (2002) [arXiv:hep-ph/0110267].
- [6] M. Alford and K. Rajagopal, JHEP **0206**, 031 (2002) [arXiv:hep-ph/0204001].
- [7] A. W. Steiner, S. Reddy and M. Prakash, Phys. Rev. D **66**, 094007 (2002) [arXiv:hep-ph/0205201].
- [8] M. Huang, P. f. Zhuang and W. q. Chao, Phys. Rev. D **67**, 065015 (2003) [arXiv:hep-ph/0207008]; F. Neumann, M. Buballa and M. Oertel, Nucl. Phys. A **714**, 481 (2003) [arXiv:hep-ph/0210078].
- [9] M. Alford, C. Kouvaris and K. Rajagopal, Phys. Rev. Lett. **92**, 222001 (2004) [arXiv:hep-ph/0311286].
- [10] M. Alford, C. Kouvaris and K. Rajagopal, Phys. Rev. D.,

- to appear, arXiv:hep-ph/0406137.
- [11] M. Alford, C. Kouvaris and K. Rajagopal, arXiv:hep-ph/0407257.
 - [12] S. B. Rüster, I. A. Shovkovy and D. H. Rischke, Nucl. Phys. A **743**, 127 (2004) [arXiv:hep-ph/0405170].
 - [13] K. Fukushima, C. Kouvaris and K. Rajagopal, Phys. Rev. D **71**, 034002 (2005) [arXiv:hep-ph/0408322].
 - [14] M. Alford, P. Jotwani, C. Kouvaris, J. Kundu and K. Rajagopal, Phys. Rev. D **71**, 114011 (2005) [arXiv:astro-ph/0411560].
 - [15] H. Abuki, M. Kitazawa and T. Kunihiro, Phys. Lett. B **615**, 102 (2005) [arXiv:hep-ph/0412382].
 - [16] S. B. Rüster, V. Werth, M. Buballa, I. A. Shovkovy and D. H. Rischke, Phys. Rev. D **72**, 034004 (2005) [arXiv:hep-ph/0503184].
 - [17] I. Shovkovy and M. Huang, Phys. Lett. B **564**, 205 (2003) [arXiv:hep-ph/0302142]; M. Huang and I. Shovkovy, arXiv:hep-ph/0307273.
 - [18] E. Gubankova, W. V. Liu and F. Wilczek, Phys. Rev. Lett. **91**, 032001 (2003).
 - [19] M. Huang and I. A. Shovkovy, Phys. Rev. D **70**, 051501 (2004) [arXiv:hep-ph/0407049]; M. Huang and I. A. Shovkovy, Phys. Rev. D **70**, 094030 (2004) [arXiv:hep-ph/0408268]; I. Giannakis and H. C. Ren, Phys. Lett. B **611**, 137 (2005) [arXiv:hep-ph/0412015]; M. Alford and Q. h. Wang, arXiv:hep-ph/0501078; M. Huang, arXiv:hep-ph/0504235.
 - [20] R. Casalbuoni, R. Gatto, M. Mannarelli, G. Nardulli and M. Ruggieri, Phys. Lett. B **605**, 362 (2005) [Erratum-ibid. B **615**, 297 (2005)] [arXiv:hep-ph/0410401]; K. Fukushima, Phys. Rev. D **72**, 074002 (2005) [arXiv:hep-ph/0506080].
 - [21] P. F. Bedaque, H. Caldas and G. Rupak, Phys. Rev. Lett. **91**, 247002 (2003) [arXiv:cond-mat/0306694]; M. M. Forbes, E. Gubankova, W. V. Liu and F. Wilczek, Phys. Rev. Lett. **94**, 017001 (2005) [arXiv:hep-ph/0405059]; J. Carlson and S. Reddy, Phys. Rev. Lett. **95**, 060401 (2005) [arXiv:cond-mat/0503256].
 - [22] M. W. Zwierlein, A. Schirotzek, C. H. Schunck and W. Ketterle, arXiv:cond-mat/0511197.
 - [23] G. B. Partridge, W. Li, R. I. Kamar, Y. Liao and R. G. Hulet, arXiv:cond-mat/0511752.
 - [24] S. Reddy and G. Rupak, Phys. Rev. C **71**, 025201 (2005) [arXiv:nucl-th/0405054].
 - [25] E. V. Gorbar, M. Hashimoto and V. A. Miransky, Phys. Lett. B **632**, 305 (2006) [arXiv:hep-ph/0507303].
 - [26] M. G. Alford, J. A. Bowers and K. Rajagopal, Phys. Rev. D **63**, 074016 (2001) [arXiv:hep-ph/0008208]; J. A. Bowers, J. Kundu, K. Rajagopal and E. Shuster, Phys. Rev. D **64**, 014024 (2001) [arXiv:hep-ph/0101067]; A. K. Leibovich, K. Rajagopal and E. Shuster, Phys. Rev. D **64**, 094005 (2001) [arXiv:hep-ph/0104073]; J. Kundu and K. Rajagopal, Phys. Rev. D **65**, 094022 (2002) [arXiv:hep-ph/0112206]; R. Casalbuoni and G. Nardulli, Rev. Mod. Phys. **263**, 320 (2004) [arXiv:hep-ph/0305069]; R. Casalbuoni, M. Ciminale, M. Mannarelli, G. Nardulli, M. Ruggieri and R. Gatto, arXiv:hep-ph/0404090.
 - [27] J. A. Bowers and K. Rajagopal, Phys. Rev. D **66**, 065002 (2002) [arXiv:hep-ph/0204079].
 - [28] R. Casalbuoni, R. Gatto, N. Ippolito, G. Nardulli and M. Ruggieri, Phys. Lett. B **627**, 89 (2005) [arXiv:hep-ph/0507247].
 - [29] A. I. Larkin and Yu. N. Ovchinnikov, Zh. Eksp. Teor. Fiz. **47**, 1136 (1964) [Sov. Phys. JETP **20**, 762 (1965)]; P. Fulde and R. A. Ferrell, Phys. Rev. **135**, A550 (1964); S. Takada and T. Izuyama, Prog. Theor. Phys. **41**, 635 (1969).
 - [30] P. F. Bedaque and T. Schäfer, Nucl. Phys. A **697**, 802 (2002) [arXiv:hep-ph/0105150]; D. B. Kaplan and S. Reddy, Phys. Rev. D **65**, 054042 (2002) [arXiv:hep-ph/0107265]; A. Kryjevski, D. B. Kaplan and T. Schäfer, Phys. Rev. D **71**, 034004 (2005) [arXiv:hep-ph/0404290]; M. Buballa, Phys. Lett. B **609**, 57 (2005) [arXiv:hep-ph/0410397]; M. M. Forbes, arXiv:hep-ph/0411001; A. Kryjevski and T. Schäfer, Phys. Lett. B **606**, 52 (2005) [arXiv:hep-ph/0407329]; A. Kryjevski and D. Yamada, Phys. Rev. D **71**, 014011 (2005) [arXiv:hep-ph/0407350].
 - [31] A. Kryjevski, arXiv:hep-ph/0508180; T. Schäfer, arXiv:hep-ph/0508190; see also D. K. Hong, arXiv:hep-ph/0506097.
 - [32] D. Bailin and A. Love, Phys. Rept. **107**, 325 (1984).
 - [33] M. G. Alford, K. Rajagopal and F. Wilczek, Phys. Lett. B **422**, 247 (1998) [arXiv:hep-ph/9711395].
 - [34] R. Rapp, T. Schäfer, E. V. Shuryak and M. Velkovsky, Phys. Rev. Lett. **81**, 53 (1998) [arXiv:hep-ph/9711396].
 - [35] M. Iwasaki and T. Iwado, Phys. Lett. B **350**, 163 (1995).
 - [36] T. Schäfer, Phys. Rev. D **62**, 094007 (2000) [arXiv:hep-ph/0006034]; M. Buballa, J. Hosek and M. Oertel, Phys. Rev. Lett. **90**, 182002 (2003) [arXiv:hep-ph/0204275]; A. Schmitt, Q. Wang and D. H. Rischke, Phys. Rev. D **66**, 114010 (2002) [arXiv:nucl-th/0209050]; M. G. Alford, J. A. Bowers, J. M. Cheyne and G. A. Cowan, Phys. Rev. D **67**, 054018 (2003) [arXiv:hep-ph/0210106]; A. Schmitt, Phys. Rev. D **71**, 054016 (2005) [arXiv:nucl-th/0412033].
 - [37] K. Rajagopal and F. Wilczek, Phys. Rev. Lett. **86**, 3492 (2001) [arXiv:hep-ph/0012039].
 - [38] A. Gerhold and A. Rebhan, Phys. Rev. D **68**, 011502 (2003) [arXiv:hep-ph/0305108]; A. Kryjevski, Phys. Rev. D **68**, 074008 (2003) [arXiv:hep-ph/0305173]; A. Gerhold, Phys. Rev. D **71**, 014039 (2005) [arXiv:hep-ph/0411086].
 - [39] D. D. Dietrich and D. H. Rischke, Prog. Part. Nucl. Phys. **53**, 305 (2004) [arXiv:nucl-th/0312044].
 - [40] F. Harary, *Graph Theory*, (Addison-Wesley, London, 1969).
 - [41] R.A. Brualdi, *Introductory Combinatorics*, (Prentice-Hall, New Jersey, 1992).
 - [42] G. Strang, *Introduction to Linear Algebra* (Wellesley-Cambridge Press, Wellesley, 1998).
 - [43] D. Vollhardt and P. Wölfle, *The Superfluid Phases of Helium 3* (Taylor & Francis, London, 1990).
 - [44] First and last items in Ref. [36].
 - [45] M. Buballa and I. A. Shovkovy, Phys. Rev. D **72**, 097501 (2005) [arXiv:hep-ph/0508197].
 - [46] K. Iida, T. Matsuura, M. Tachibana and T. Hatsuda, Phys. Rev. Lett. **93**, 132001 (2004) [arXiv:hep-ph/0312363].
 - [47] A. Schmitt, *Ph.D. dissertation*, arXiv:nucl-th/0405076.
 - [48] H. Abuki and T. Kunihiro, arXiv:hep-ph/0509172.